

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>A01H 5/00, 1/02, C09K 7/02, C12P 7/40, 7/64, C12N 9/20</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/07433</b> <b>(43) International Publication Date:</b> 17 February 2000 (17.02.00)
<b>(21) International Application Number:</b> PCT/US99/17645 <b>(22) International Filing Date:</b> 3 August 1999 (03.08.99) <b>(30) Priority Data:</b> 09/128,602 3 August 1998 (03.08.98) US <b>(71) Applicant (for all designated States except US):</b> CARGILL, INCORPORATED [US/US]; 15407 West McGinty Road, Wayzata, MN 55441-5624 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> KODALI, Dharma [IN/US]; 710 Olive Lane North, Plymouth, MN 55447 (US). FAN, Zhegong [CN/US]; 4429 Monte Carlo Place, Fort Collins, CO 80525 (US). DeBONTE, Lorin, R. [US/US]; 3706 Rochdale Drive, Fort Collins, CO 80525 (US). <b>(74) Agents:</b> MCCORMICK GRAHAM, Monica et al.; Fish & Richardson P.C., P.A., Suite 3300, 60 South Sixth Street, Minneapolis, MN 55402 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> PLANTS, SEEDS AND OILS HAVING AN ELEVATED TOTAL MONOUNSATURATED FATTY ACID CONTENT		
<b>(57) Abstract</b>  Plants, seeds and oils having a total long-chain monounsaturated content of at least about 82 % and an erucic acid content of at least about 15 % are described. Methods for producing plants having the profiled fatty acid content are also described.		

77  
15  
21  
4  
117

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5

Plants, Seeds and Oils Having an Elevated Total  
Monounsaturated Fatty Acid Content

Technical Field

This invention relates to fatty acid desaturases and nucleic acids encoding desaturase proteins. More particularly, the invention relates to nucleic acids encoding delta-12 and delta-15 fatty acid desaturase proteins that affect fatty acid composition in plants, polypeptides produced from such nucleic acids and plants expressing such nucleic acids.

Background of the Invention

Many breeding studies have been conducted to improve the fatty acid profile of *Brassica* varieties. Pleines and Freidt, *Fat Sci. Technol.*, 90(5), 167-171 (1988) report plant lines with reduced  $C_{18:3}$  levels (2.5-5.8%) combined with high oleic content (73-79%). Rakow and McGregor, *J. Amer. Oil Chem. Soc.*, 50, 400-403 (Oct. 1973) discuss problems associated with selecting mutants for linoleic and linolenic acids. In *Can. J. Plant Sci.*, 68, 509-511 (Apr. 1988) Stellar summer rape producing seed oil with 3% linolenic acid and 28% linoleic acid is disclosed. Roy and Tarr, *Z. Pflanzenzuchtg.*, 95(3), 201-209 (1985) report transfer of genes through an interspecific cross from *Brassica juncea* into *Brassica napus* resulting in a reconstituted line combining high linoleic with low linolenic acid content. Roy and Tarr, *Plant Breeding*, 98, 89-96 (1987) discuss prospects for development of *B. napus* L. having improved linolenic and linolenic acid content. European Patent application 323,753 published July 12, 1989 discloses seeds and oils having greater than 79% oleic acid combined with less than 3.5% linolenic acid. Canvin, *Can. J. Botany*, 43, 63-69 (1965) discusses the effect of temperature on the fatty acid composition of oils from several seed crops including rapeseed.

Mutations typically are induced with extremely high doses of radiation and/or chemical mutagens (Gaul, *H. Radiation Botany* (1964) 4:155-232). High

- 2 -

dose levels which exceed LD<sub>50</sub>, and typically reach LD<sub>90</sub>, led to maximum achievable mutation rates. In mutation breeding of *Brassica* varieties, high levels of chemical mutagens alone or combined with radiation have induced a limited number of fatty acid mutations (Rakow, G.Z. *Pflanzenzuchtg* (1973) 69:62-82).

- 5 The low  $\alpha$ -linolenic acid mutation derived from the Rakow mutation breeding program did not have direct commercial application because of low seed yield. The first commercial cultivar using the low  $\alpha$ -linolenic acid mutation derived in 1973 was released in 1988 as the variety Stellar (Scarth, R. et al., *Can. J. Plant Sci.* (1988) 68:509-511). Stellar was 20% lower yielding than commercial cultivars at  
10 the time of its release.

Alterations in fatty acid composition of vegetable oils is desirable for meeting specific food and industrial uses. For example, *Brassica* canola varieties with increased monounsaturate levels (oleic acid) in the seed oil, and products derived from such oil, would improve lipid nutrition. Canola lines which are low  
15 in polyunsaturated fatty acids and high in oleic acid tend to have higher oxidative stability, which is a useful trait for the retail food industry. Useful traits of vegetable oils for industrial uses like lubrication fluids include desirable low temperature behavior such as low pour point and low cloud point along with very high oxidative stability.

- 20 Delta-12 fatty acid desaturase (also known as oleic desaturase) is involved in the enzymatic conversion of oleic acid to linoleic acid. Delta-15 fatty acid desaturase (also known as linoleic acid desaturase) is involved in the enzymatic conversion of linoleic acid to  $\alpha$ -linolenic acid. A microsomal delta-12 desaturase has been cloned and characterized using T-DNA tagging. Okuley, et al.,  
25 *Plant Cell* 6:147-158 (1994). The nucleotide sequences of higher plant genes encoding microsomal delta-12 fatty acid desaturase are described in Lightner et al., WO94/11516. Sequences of higher plant genes encoding microsomal and plastid delta-15 fatty acid desaturases are disclosed in Yadav, N., et al., *Plant Physiol.*, 103:467-476 (1993), WO 93/11245 and Arondel, V. et al., *Science*, 258:1353-1355  
30 (1992).



- 3 -

Summary of the Invention

Triacylglycerols containing fatty acids with heterogenous chain lengths and with high monounsaturate levels can provide useful traits for industrial purposes. Plants with fatty acid compositions that have high monounsaturate levels  
5 and heterogenous chain lengths would provide a source of industrial oils for uses such as lubrication.

In one aspect, the invention features a *Brassica* plant, and progeny thereof, producing seeds having a long chain monounsaturated fatty acid content of at least about 82% and an erucic acid content of at least about 15% based on total  
10 fatty acid composition. The oleic acid and eicosenoic acid content of the seeds is at least about 37% and at least about 14%, based on total fatty acid composition, respectively. The saturated fatty acid content of such seeds is less than 7% and the polyunsaturated fatty acid content is less than about 11%.

In some embodiments, the plants have a monounsaturated fatty acid  
15 content of from about 85% to about 90% and an erucic acid content of at least about 15% based on total fatty acid composition. In such plants, the oleic acid content can be at least about 42% and in particular, from about 47% to about 56% based on total fatty acid composition. The erucic acid content is from about 17% to about 31%, and the eicosenoic acid content is from about 15% to about 21%.

20 The invention also features a *Brassica* seed oil having a long chain monounsaturated fatty acid content of at least about 82% and an erucic acid content of at least about 15% based on total fatty acid composition. Such oils can have an oleic acid and eicosenoic acid content of at least about 14% and 37%, respectively, based on total fatty acid composition. The saturated fatty acid content can be less  
25 than about 7%, e.g., less than about 4% or about 2 to 4%. The polyunsaturated fatty acid content is less than about 11% and in particular embodiments, less than 9%, based on total fatty acid composition. The  $\alpha$ -linolenic acid content can be about 1% to about 2%.

In some embodiments, the *Brassica* seed oil contains a long chain  
30 monounsaturated fatty acid content of from about 85% to about 90%. In such oils,

- 4 -

the oleic acid content is at least about 42%, and in particular embodiments, is from about 47% to about 56%, based on total fatty acid composition. The erucic acid and eicosenoic acid content is from about 17% to about 31% and from about 15% to about 21%, respectively, based on total fatty acid composition. Also featured is

5 a *Brassica* seed oil having a long chain monounsaturated fatty acid content of at least about 82%, wherein the sum of the nervonic acid, erucic acid and eicosenoic acid content is from about 50% to about 66% based on total fatty acid composition. Such a seed oil can have an oleic acid content from about 25% to about 30%.

The invention also features a method of producing plants having a long

10 chain monounsaturated fatty acid content of at least about 82% and an erucic acid content of at least about 15%, based on total fatty acid composition. The methods include crossing a first plant line with a second plant line and selecting progeny with the desired fatty acid composition. The first plant line has an erucic acid content of at least about 45%. The second plant line has an oleic acid content of at

15 least about 84%.

The invention also features a method of making a vegetable oil. The method comprises the steps of crushing *Brassica* seeds having a long chain monounsaturated fatty acid content of at least about 82% and an erucic acid content of at least about 15% based on total fatty acid composition, and extracting the

20 vegetable oil from the crushed seeds. The method can also include the steps of refining and bleaching the oil, and deodorizing the oil.

The invention also features a lubricant or hydraulic fluid comprising a *Brassica* oil having a long chain monounsaturated fatty acid content of at least about 82% and an erucic acid content of at least about 15% based on total fatty

25 acid composition, and an additive. The additive can be an antioxidant, a rust inhibitor, a corrosion inhibitor, a pour point depressant, an anti-foam additive, a colorant and a detergent. The additive can be present in an amount from about 0.01% to about 20% by weight.

- 5 -

### Brief Description of the Sequence Listing

SEQ ID NO:1 shows the DNA sequence for the coding region of a wild type *Brassica* Fad2-D gene. SEQ ID NO:2 is the deduced amino acid sequence for SEQ ID NO:1.

5        SEQ ID NO:3 shows the DNA sequence for the coding region of the IMC 129 mutant *Brassica* Fad2-D gene. SEQ ID NO:4 is the deduced amino acid sequence for SEQ ID NO:3.

      SEQ ID NO:5 shows the DNA sequence for the coding region of a wild type *Brassica* Fad2-F gene. SEQ ID NO:6 is the deduced amino acid sequence for  
10    SEQ ID NO:5.

      SEQ ID NO:7 shows the DNA sequence for the coding region of the Q508 mutant *Brassica* Fad2-F gene. SEQ ID NO:8 is the deduced amino acid sequence for SEQ ID NO:7.

      SEQ ID NO:9 shows the DNA sequence for the coding region of the  
15    Q4275 mutant *Brassica* Fad2-F gene. SEQ ID NO:10 is the deduced amino acid sequence for SEQ ID NO:9.

### Brief Description of the Figures

Figure 1 is a histogram showing the frequency distribution of seed oil oleic acid ( $C_{18:1}$ ) content in a segregating population of a Q508 X Westar cross.

20    The bar labeled WSGA 1A represents the  $C_{18:1}$  content of the Westar parent. The bar labeled Q508 represents the  $C_{18:1}$  content of the Q508 parent.

      Figure 2 shows the nucleotide sequences for a *Brassica* Fad2-D wild type gene (Fad2-D wt), IMC129 mutant gene (Fad2-D GA316 IMC129), Fad2-F wild type gene (Fad2-F wt), Q508 mutant gene (Fad2-F TA515 Q508) and Q4275  
25    mutant gene (Fad2-F GA908 Q4275).

      Figure 3 shows the deduced amino acid sequences for the polynucleotides of Figure 2.

      Figure 4 is a schematic of a breeding procedure used to produce *Brassica* plants having a high erucic acid and a high oleic acid content.

- 6 -

Detailed Description

All percent fatty acids herein are percent by weight of the oil of which the fatty acid is a component.

As used herein, a "line" is a group of plants that display little or no genetic variation between individuals for at least one trait. Such lines may be created by several generations of self-pollination and selection, or vegetative propagation from a single parent using tissue or cell culture techniques. As used herein, the term "variety" refers to a line which is used for commercial production.

The term "mutagenesis" refers to the use of a mutagenic agent to induce random genetic mutations within a population of individuals. The treated population, or a subsequent generation of that population, is then screened for usable trait(s) that result from the mutations. A "population" is any group of individuals that share a common gene pool. As used herein "M<sub>0</sub>" is untreated seed. As used herein, "M<sub>1</sub>" is the seed (and resulting plants) exposed to a mutagenic agent, while "M<sub>2</sub>" is the progeny (seeds and plants) of self-pollinated M<sub>1</sub> plants, "M<sub>3</sub>" is the progeny of self-pollinated M<sub>2</sub> plants, and "M<sub>4</sub>" is the progeny of self-pollinated M<sub>3</sub> plants. "M<sub>5</sub>" is the progeny of self-pollinated M<sub>4</sub> plants. "M<sub>6</sub>", "M<sub>7</sub>", etc. are each the progeny of self-pollinated plants of the previous generation. The term "selfed" as used herein means self-pollinated.

"Stability" or "stable" as used herein means that with respect to a given fatty acid component, the component is maintained from generation to generation for at least two generations and preferably at least three generations at substantially the same level, e.g., preferably  $\pm 5\%$ . The method of invention is capable of creating lines with improved fatty acid compositions stable up to  $\pm 5\%$  from generation to generation. The above stability may be affected by temperature, location, stress and time of planting. Thus, comparison of fatty acid profiles should be made from seeds produced under similar growing conditions. Stability may be measured based on knowledge of prior generation.

Intensive breeding has produced certain *Brassica* plants whose seed oil contains less than 2% erucic acid. The same varieties have also been bred so that

- 7 -

the defatted meal contains less than 30  $\mu\text{mol}$  glucosinolates/gram. "Canola" as used herein refers to plant seeds or oils which contain less than 2% erucic acid ( $\text{C}_{22:1}$ ), and result in a defatted meal with less than 30  $\mu\text{mol}$  glucosinolates/gram.

Applicants have discovered plants with mutations in a delta-12 fatty acid  
5 desaturase gene. Such plants have useful alterations in the fatty acid compositions of the seed oil. Such mutations confer, for example, an elevated oleic acid content, a decreased, stabilized linoleic acid content, or both elevated oleic acid and decreased, stabilized linoleic acid content.

Applicants have further discovered plants with mutations in a delta-15  
10 fatty acid desaturase gene. Such plants have useful alterations in the fatty acid composition of the seed oil, e.g., a decreased, stabilized level of  $\alpha$ -linolenic acid.

Applicants have further discovered isolated nucleic acid fragments (polynucleotides) comprising sequences that carry mutations within the coding sequence of delta-12 or delta-15 fatty acid desaturases. The mutations confer  
15 desirable alterations in fatty acid levels in the seed oil of plants carrying such mutations. Delta-12 fatty acid desaturase is also known as omega-6 fatty acid desaturase and is sometimes referred to herein as Fad2 or 12-DES. Delta-15 fatty acid desaturase is also known as omega-3 fatty acid desaturase and is sometimes referred to herein as Fad3 or 15-DES.

20 A nucleic acid fragment of the invention may be in the form of RNA or in the form of DNA, including cDNA, synthetic DNA or genomic DNA. The DNA may be double-stranded or single-stranded, and if single-stranded, can be either the coding strand or non-coding strand. An RNA analog may be, for example, mRNA or a combination of ribo- and deoxyribonucleotides. Illustrative  
25 examples of a nucleic acid fragment of the invention are the mutant sequences shown in Fig. 3.

A nucleic acid fragment of the invention contains a mutation in a microsomal delta-12 fatty acid desaturase coding sequence or a mutation in a microsomal delta-15 fatty acid desaturase coding sequence. Such a mutation  
30 renders the resulting desaturase gene product non-functional in plants, relative to

- 8 -

the function of the gene product encoded by the wild-type sequence. The non-functionality of the delta-12 desaturase gene product can be inferred from the decreased level of reaction product (linoleic acid) and increased level of substrate (oleic acid) in plant tissues expressing the mutant sequence, compared to the  
5 corresponding levels in plant tissues expressing the wild-type sequence. The non-functionality of the delta-15 desaturase gene product can be inferred from the decreased level of reaction product ( $\alpha$ -linolenic acid) and the increased level of substrate (linoleic acid) in plant tissues expressing the mutant sequence, compared to the corresponding levels in plant tissues expressing the wild-type sequence.

10 A nucleic acid fragment of the invention may comprise a portion of the coding sequence, e.g., at least about 10 nucleotides, provided that the fragment contains at least one mutation in the coding sequence. The length of a desired fragment depends upon the purpose for which the fragment will be used, e.g., PCR primer, site-directed mutagenesis and the like. In one embodiment, a nucleic acid  
15 fragment of the invention comprises the full length coding sequence of a mutant delta-12 or mutant delta-15 fatty acid desaturase, e.g., the mutant sequences of Fig. 3. In other embodiments, a nucleic acid fragment is about 20 to about 50 nucleotides (or base pairs, bp), or about 50 to about 500 nucleotides, or about 500 to about 1200 nucleotides in length.

20 Desirable alterations in fatty acid levels in the seed oil of plants can be produced using a ribozyme. Ribozyme molecules designed to cleave delta-12 or delta-15 desaturase mRNA transcripts can be used to prevent expression of delta-12 or delta-15 desaturases. While various ribozymes that cleave mRNA at site-specific recognition sequences can be used to destroy desaturase mRNAs, hammerhead  
25 ribozymes are particularly useful. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole requirement is that the target RNA contain a 5'-UG-3' nucleotide sequence. The construction and production of hammerhead ribozymes is well known in the art. See, for example, U.S. Patent No. 5,254,678. Hammerhead  
30 ribozyme sequences can be embedded in a stable RNA such as a transfer RNA

- 9 -

(tRNA) to increase cleavage efficiency *in vivo*. Perriman, R. et al., Proc. Natl. Acad. Sci. USA, 92(13):6175-6179 (1995); de Feyter, R. and Gaudron, J., Methods in Molecular Biology, Vol. 74, Chapter 43, "Expressing Ribozymes in Plants", Edited by Turner, P.C, Humana Press Inc., Totowa, NJ. RNA endoribonucleases  
5 such as the one that occurs naturally in *Tetrahymena thermophila*, and which have been described extensively by Cech and collaborators are also useful. See, for example, U.S. Patent No. 4,987,071.

A mutation in a nucleic acid fragment of the invention may be in any portion of the coding sequence that renders the resulting gene product non-  
10 functional. Suitable types of mutations include, without limitation, insertions of nucleotides, deletions of nucleotides, or transitions and transversions in the wild-type coding sequence. Such mutations result in insertions of one or more amino acids, deletions of one or more amino acids, and non-conservative amino acid substitutions in the corresponding gene product. In some embodiments, the  
15 sequence of a nucleic acid fragment may comprise more than one mutation or more than one type of mutation.

Insertion or deletion of amino acids in a coding sequence may, for example, disrupt the conformation of essential alpha-helical or beta-pleated sheet regions of the resulting gene product. Amino acid insertions or deletions may also  
20 disrupt binding or catalytic sites important for gene product activity. It is known in the art that the insertion or deletion of a larger number of contiguous amino acids is more likely to render the gene product non-functional, compared to a smaller number of inserted or deleted amino acids. Non-conservative amino acid substitutions may replace an amino acid of one class with an amino acid of a  
25 different class. Non-conservative substitutions may make a substantial change in the charge or hydrophobicity of the gene product. Non-conservative amino acid substitutions may also make a substantial change in the bulk of the residue side chain, e.g., substituting an alanyl residue for a isoleucyl residue.

Examples of non-conservative substitutions include the substitution of a  
30 basic amino acid for a non-polar amino acid, or a polar amino acid for an acidic

- 10 -

amino acid. Because there are only 20 amino acids encoded in a gene, substitutions that result in a non-functional gene product may be determined by routine experimentation, incorporating amino acids of a different class in the region of the gene product targeted for mutation.

5 Preferred mutations are in a region of the nucleic acid encoding an amino acid sequence motif that is conserved among delta-12 fatty acid desaturases or delta-15 fatty acid desaturases, such as a His-Xaa-Xaa-Xaa-His motif (Tables 1-3). An example of a suitable region has a conserved HECGH motif that is found, for example, in nucleotides corresponding to amino acids 105 to 109 of the  
10 *Arabidopsis* and *Brassica* delta-12 desaturase sequences, in nucleotides corresponding to amino acids 101 to 105 of the soybean delta-12 desaturase sequence and in nucleotides corresponding to amino acids 111 to 115 of the maize delta-12 desaturase sequence. See e.g., WO 94/115116; Okuley et al., Plant Cell 6:147-158 (1994). The one letter amino acid designations used herein are  
15 described in Alberts, B. et al., Molecular Biology of the Cell, 3rd edition, Garland Publishing, New York, 1994. Amino acids flanking this motif are also highly conserved among delta-12 and delta-15 desaturases and are also suitable candidates for mutations in fragments of the invention.

An illustrative embodiment of a mutation in a nucleic acid fragment of  
20 the invention is a Glu to Lys substitution in the HECGH motif of a *Brassica* microsomal delta-12 desaturase sequence, either the D form or the F form. This mutation results in the sequence HECGH being changed to HKCGH as seen by comparing SEQ ID NO:2 (wild-type D form) to SEQ ID NO:4 (mutant D form). A similar mutation in other Fad-2 sequences is contemplated to result in a non-  
25 functional gene product.

A similar motif may be found at amino acids 101 to 105 of the *Arabidopsis* microsomal delta-15 fatty acid desaturase, as well as in the corresponding rape and soybean desaturases (Table 5). See, e.g., WO 93/11245; Arondel, V. et al., Science, 258:1153-1155 (1992); Yadav, N. et al., Plant Physiol.,  
30 103:467-476 (1993). Plastid delta-15 fatty acids have a similar motif (Table 5).



- 11 -

Among the types of mutations in an HECGH motif that render the resulting gene product non-functional are non-conservative substitutions. An illustrative example of a non-conservative substitution is substitution of a glycine residue for either the first or second histidine. Such a substitution replaces a

5 charged residue (histidine) with a non-polar residue (glycine). Another type of mutation that renders the resulting gene product non-functional is an insertion mutation, e.g., insertion of a glycine between the cysteine and glutamic acid residues in the HECGH motif.

Other regions having suitable conserved amino acid motifs include the

10 HRRHH motif shown in Table 2, the HRTTH motif shown in Table 6 and the HVAHH motif shown in Table 3. See, e.g., WO 94/115116; Hitz, W. et al., Plant Physiol., 105:635-641 (1994); Okuley, J., et al., supra; and Yadav, N. et al., supra. An illustrative example of a mutation in the region shown in Table 3 is a mutation at nucleotides corresponding to the codon for glycine (amino acid 303 of *B. napus*).

15 A non-conservative Gly to Glu substitution results in the amino acid sequence DRDYGILNKV being changed to sequence DRDYEILNKV (compare wild-type F form SEQ ID NO:6 to mutant Q4275 SEQ ID NO:10, Fig. 3).

Another region suitable for a mutation in a delta-12 desaturase sequence contains the motif KYLNNP at nucleotides corresponding to amino acids 171 to

20 175 of the *Brassica* desaturase sequence. An illustrative example of a mutation in this region is a Leu to His substitution, resulting in the amino acid sequence (Table 4) KYHNN (compare wild-type Fad2-F SEQ ID NO:6 to mutant SEQ ID NO:8). A similar mutation in other Fad-2 amino acid sequences is contemplated to result in a non-functional gene product.

- 12 -

TABLE 1

Alignment of Amino Acid Sequences from Microsomal  
Delta-12 Fatty Acid Desaturases

	<u>Species</u>	<u>Position</u>	<u>Amino Acid Sequence</u>
5	<i>Arabidopsis thaliana</i>	100-129	IWVIAHECGH HAFSDYQWLD DTVGLIFHSF
	<i>Glycine max</i>	96-125	VWVIAHECGH HAFSKYQWVD DVVGLTLHST
	<i>Zea mays</i>	106-135	VWVIAHECGH HAFSDYSLLD DVVGLVLHSS
	<i>Ricinus communis</i> <sup>a</sup>	1- 29	WVMAHDCGH HAFSDYQLLD DVVGLILHSC
	<i>Brassica napus D</i>	100-128	VWVIAHECGH HAFSDYQWLD DTVGLIFHS
10	<i>Brassica napus F</i>	100-128	VWVIAHECGH HAFSDYQWLD DTVGLIFHS

<sup>a</sup> from plasmid pRF2-1C

TABLE 2

Alignment of Amino Acid Sequences from Microsomal  
Delta-12 Fatty Acid Desaturases

	<u>Species</u>	<u>Position</u>	<u>Amino Acid Sequence</u>
15	<i>Arabidopsis thaliana</i>	130-158	LLVPYFSWKY SHRRHHSNTG SLERDEVFV
	<i>Glycine max</i>	126-154	LLVPYFSWKI SHRRHHSNTG SLDRDEVFV
	<i>Zea mays</i>	136-164	LMVPYFSWKY SHRRHHSNTG SLERDEVFV
	<i>Ricinus communis</i> <sup>a</sup>	30- 58	LLVPYFSWKH SHRRHHSNTG SLERDEVFV
20	<i>Brassica napus D</i>	130-158	LLVPYFSWKY SHRRHHSNTG SLERDEVFV
	<i>Brassica napus F</i>	130-158	LLVPYFSWKY SHRRHHSNTG SLERDEVFV

<sup>a</sup> from plasmid pRF2-1C

TABLE 3

Alignment of Amino Acid Sequences from Microsomal  
Delta-12 Fatty Acid Desaturases

	<u>Species</u>	<u>Position</u>	<u>Amino Acid Sequence</u>
25	<i>Arabidopsis thaliana</i>	298-333	DRDYGILNKV FHNITDTHVA HHLFSTMPHY NAMEAT
	<i>Glycine max</i>	294-329	DRDYGILNKV FHNITDTHVA HHLFSTMPHY HAMEAT
	<i>Zea mays</i>	305-340	DRDYGILNRV FHNITDTHVA HHLFSTMPHY HAMEAT
30	<i>Ricinus communis</i> <sup>a</sup>	198-224	DRDYGILNKV FHNITDTQVA HHLF TMP
	<i>Brassica napus D</i>	299-334	DRDYGILNKV FHNITDTHVA HHLFSTMPHY HAMEAT
	<i>Brassica napus F</i>	299-334	DRDYGILNKV FHNITDTHVA HHLFSTMPHY HAMEAT

<sup>a</sup> from plasmid pRF2-1C

- 13 -

**TABLE 4**

Alignment of Conserved Amino Acids from Microsomal  
Delta-12 Fatty Acid Desaturases

	<u>Species</u>	<u>Position</u>	<u>Amino Acid Sequence</u>
5	<i>Arabidopsis thaliana</i>	165-180	IKWYGKYLNN PLGRIM
	<i>Glycine max</i>	161-176	VAWFSLYLNN PLGRAV
	<i>Zea mays</i>	172-187	PWYTPYVYNN PVGRVV
	<i>Ricinus communis</i> <sup>a</sup>	65- 80	IRWYSKYLNN PPGRIM
	<i>Brassica napus</i> D	165-180	IKWYGKYLNN PLGRTV
10	<i>Brassica napus</i> F	165-180	IKWYGKYLNN PLGRTV
	<sup>a</sup> from plasmid pRF2-1C		

**TABLE 5**

Alignment of Conserved Amino Acids from Plastid and Microsomal  
Delta-15 Fatty Acid Desaturases

	<u>Species</u>	<u>Position</u>	<u>Amino Acid Sequence</u>
15	<i>Arabidopsis thaliana</i> <sup>a</sup>	156-177	WALFVLGHD CGHGSFSNDP KLN
	<i>Brassica napus</i> <sup>a</sup>	114-135	WALFVLGHD CGHGSFSNDP RLN
	<i>Glycine max</i> <sup>a</sup>	164-185	WALFVLGHD CGHGSFSNNS KLN
	<i>Arabidopsis thaliana</i>	94-115	WAIFVLGHD CGHGSFSDIP LLN
20	<i>Brassica napus</i>	87-109	WALFVLGHD CGHGSFSNDP RLN
	<i>Glycine max</i>	93-114	WALFVLGHD CGHGSFSDSP PLN

<sup>a</sup> Plastid sequences

- 14 -

**TABLE 6**  
Alignment of Conserved Amino Acids from Plastid and Microsomal  
Delta-15 Fatty Acid Desaturases

	<u>Species</u>	<u>Position</u>	<u>Amino Acid Sequence</u>
5	<i>A. thaliana</i> <sup>a</sup>	188-216	ILVPYHGWRI SHRTHHQNHG HVENDESWH
	<i>B. napus</i> <sup>a</sup>	146-174	ILVPYHGWRI SHRTHHQNHG HVENDESWH
	<i>Glycine max</i> <sup>a</sup>	196-224	ILVPYHGWRI SHRTHHQHHG HAENDESWH
	<i>A. thaliana</i>	126-154	ILVPYHGWRI SHRTHHQNHG HVENDESWV
	<i>Brassica napus</i>	117-145	ILVPYHGWRI SHRTHHQNHG HVENDESWV
10	<i>Glycine max</i>	125-153	ILVPYHGWRI SHRTHHQNHG HIEKDESWV

<sup>a</sup> Plastid sequences

The conservation of amino acid motifs and their relative positions indicates that regions of a delta-12 or delta-15 fatty acid desaturase that can be mutated in one species to generate a non-functional desaturase can be mutated in the corresponding region from other species to generate a non-functional delta-12 desaturase or delta-15 desaturase gene product in that species.

Mutations in any of the regions of Tables 1-6 are specifically included within the scope of the invention and are substantially identical to those mutations exemplified herein, provided that such mutation (or mutations) renders the resulting desaturase gene product non-functional, as discussed hereinabove.

A nucleic acid fragment containing a mutant sequence can be generated by techniques known to the skilled artisan. Such techniques include, without limitation, site-directed mutagenesis of wild-type sequences and direct synthesis using automated DNA synthesizers.

A nucleic acid fragment containing a mutant sequence can also be generated by mutagenesis of plant seeds or regenerable plant tissue by, e.g., ethyl methane sulfonate, X-rays or other mutagens. With mutagenesis, mutant plants having the desired fatty acid phenotype in seeds are identified by known techniques and a nucleic acid fragment containing the desired mutation is isolated from genomic DNA or RNA of the mutant line. The site of the specific mutation is then

- 15 -

determined by sequencing the coding region of the delta-12 desaturase or delta-15 desaturase gene. Alternatively, labeled nucleic acid probes that are specific for desired mutational events can be used to rapidly screen a mutagenized population.

The disclosed method may be applied to all oilseed *Brassica* species,  
5 and to both Spring and Winter maturing types within each species. Physical mutagens, including but not limited to X-rays, UV rays, and other physical treatments which cause chromosome damage, and other chemical mutagens, including but not limited to ethidium bromide, nitrosoguanidine, diepoxybutane etc. may also be used to induce mutations. The mutagenesis treatment may also be  
10 applied to other stages of plant development, including but not limited to cell cultures, embryos, microspores and shoot apices.

"Stable mutations" as used herein are defined as  $M_5$  or more advanced lines which maintain a selected altered fatty acid profile for a minimum of three generations, including a minimum of two generations under field conditions, and  
15 exceeding established statistical thresholds for a minimum of two generations, as determined by gas chromatographic analysis of a minimum of 10 randomly selected seeds bulked together. Alternatively, stability may be measured in the same way by comparing to subsequent generations. In subsequent generations, stability is defined as having similar fatty acid profiles in the seed as that of the prior or  
20 subsequent generation when grown under substantially similar conditions.

Mutation breeding has traditionally produced plants carrying, in addition to the trait of interest, multiple, deleterious traits, e.g., reduced plant vigor and reduced fertility. Such traits may indirectly affect fatty acid composition, producing an unstable mutation; and/or reduce yield, thereby reducing the  
25 commercial utility of the invention. To eliminate the occurrence of deleterious mutations and reduce the load of mutations carried by the plant, a low mutagen dose is used in the seed treatments to create an LD30 population. This allows for the rapid selection of single gene mutations for fatty acid traits in agronomic backgrounds which produce acceptable yields.

- 16 -

The seeds of several different fatty acid lines have been deposited with the American Type Culture Collection and have the following accession numbers.

<u>Line</u>	<u>Accession No.</u>	<u>Deposit Date</u>
A129.5	40811	May 25, 1990
5 A133.1	40812	May 25, 1990
M3032.1	75021	June 7, 1991
M3062.8	75025	June 7, 1991
M3028.10	75026	June 7, 1991
IMC130	75446	April 16, 1993
10 Q4275	97569	May 10, 1996

In some plant species or varieties more than one form of endogenous microsomal delta-12 desaturase may be found. In amphidiploids, each form may be derived from one of the parent genomes making up the species under consideration. Plants with mutations in both forms have a fatty acid profile that differs from plants with a mutation in only one form. An example of such a plant is *Brassica napus* line Q508, a doubly-mutagenized line containing a mutant D-form of delta-12 desaturase (SEQ ID NO:3) and a mutant F-form of delta-12 desaturase (SEQ ID NO:7). Another example is line Q4275, which contains a mutant D-form of delta-12 desaturase (SEQ ID NO:3) and a mutant F-form of delta-12 desaturase (SEQ ID NO:9). See Figs. 2-3.

Preferred host or recipient organisms for introduction of a nucleic acid fragment of the invention are the oil-producing species, such as soybean (*Glycine max*), rapeseed (e.g., *Brassica napus*, *B. rapa* and *B. juncea*), sunflower (*Helianthus annuus*), castor bean (*Ricinus communis*), corn (*Zea mays*), and safflower (*Carthamus tinctorius*).

A nucleic acid fragment of the invention may further comprise additional nucleic acids. For example, a nucleic acid encoding a secretory or leader amino acid sequence can be linked to a mutant desaturase nucleic acid fragment such that the secretory or leader sequence is fused in-frame to the amino terminal end of a mutant delta-12 or delta-15 desaturase polypeptide. Other nucleic acid

- 17 -

fragments are known in the art that encode amino acid sequences useful for fusing in-frame to the mutant desaturase polypeptides disclosed herein. See, e.g., U.S. 5,629,193 incorporated herein by reference. A nucleic acid fragment may also have one or more regulatory elements operably linked thereto.

5           The present invention also comprises nucleic acid fragments that selectively hybridize to mutant desaturase sequences. Such a nucleic acid fragment typically is at least 15 nucleotides in length. Hybridization typically involves Southern analysis (Southern blotting), a method by which the presence of DNA sequences in a target nucleic acid mixture are identified by hybridization to a  
10   labeled oligonucleotide or DNA fragment probe. Southern analysis typically involves electrophoretic separation of DNA digests on agarose gels, denaturation of the DNA after electrophoretic separation, and transfer of the DNA to nitrocellulose, nylon, or another suitable membrane support for analysis with a radiolabeled, biotinylated, or enzyme-labeled probe as described in sections 9.37-9.52 of  
15   Sambrook et al., (1989) *Molecular Cloning*, second edition, Cold Spring Harbor Laboratory, Plainview; NY.

A nucleic acid fragment can hybridize under moderate stringency conditions or, preferably, under high stringency conditions to a mutant desaturase sequence. High stringency conditions are used to identify nucleic acids that have a  
20   high degree of homology to the probe. High stringency conditions can include the use of low ionic strength and high temperature for washing, for example, 0.015 M NaCl/0.0015 M sodium citrate (0.1X SSC); 0.1% sodium lauryl sulfate (SDS) at 50-65°C. Alternatively, a denaturing agent such as formamide can be employed during hybridization, e.g., 50% formamide with 0.1% bovine serum albumin/0.1%  
25   Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 with 750 mM NaCl, 75 mM sodium citrate at 42°C. Another example is the use of 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at  
30   42°C, with washes at 42°C in 0.2 x SSC and 0.1% SDS.

- 18 -

Moderate stringency conditions refers to hybridization conditions used to identify nucleic acids that have a lower degree of identity to the probe than do nucleic acids identified under high stringency conditions. Moderate stringency conditions can include the use of higher ionic strength and/or lower temperatures for washing of the hybridization membrane, compared to the ionic strength and temperatures used for high stringency hybridization. For example, a wash solution comprising 0.060 M NaCl/0.0060 M sodium citrate (4X SSC) and 0.1% sodium lauryl sulfate (SDS) can be used at 50°C, with a last wash in 1X SSC, at 65°C. Alternatively, a hybridization wash in 1X SSC at 37°C can be used.

Hybridization can also be done by Northern analysis (Northern blotting), a method used to identify RNAs that hybridize to a known probe such as an oligonucleotide, DNA fragment, cDNA or fragment thereof, or RNA fragment. The probe is labeled with a radioisotope such as  $^{32}\text{P}$ , by biotinylation or with an enzyme. The RNA to be analyzed can be usually electrophoretically separated on an agarose or polyacrylamide gel, transferred to nitrocellulose, nylon, or other suitable membrane, and hybridized with the probe, using standard techniques well known in the art such as those described in sections 7.39-7.52 of Sambrook et al., *supra*.

A polypeptide of the invention comprises an isolated polypeptide having a mutant amino acid sequence, as well as derivatives and analogs thereof. See, e.g., the mutant amino acid sequences of Fig. 3. By "isolated" is meant a polypeptide that is expressed and produced in an environment other than the environment in which the polypeptide is naturally expressed and produced. For example, a plant polypeptide is isolated when expressed and produced in bacteria or fungi. A polypeptide of the invention also comprises variants of the mutant desaturase polypeptides disclosed herein, as discussed above.

In one embodiment of the claimed invention, a plant contains both a delta-12 desaturase mutation and a delta-15 desaturase mutation. Such plants can have a fatty acid composition comprising very high oleic acid and very low alpha-linolenic acid levels. Mutations in delta-12 desaturase and delta-15 desaturase may



- 19 -

be combined in a plant by making a genetic cross between delta-12 desaturase and delta-15 desaturase single mutant lines. A plant having a mutation in delta-12 fatty acid desaturase is crossed or mated with a second plant having a mutation in delta-15 fatty acid desaturase. Seeds produced from the cross are planted and the  
5 resulting plants are selfed in order to obtain progeny seeds. These progeny seeds are then screened in order to identify those seeds carrying both mutant genes.

Alternatively, a line possessing either a delta-12 desaturase or a delta-15 desaturase mutation can be subjected to mutagenesis to generate a plant or plant line having mutations in both delta-12 desaturase and delta-15 desaturase. For  
10 example, the IMC 129 line has a mutation in the coding region (Glu<sub>106</sub> to Lys<sub>106</sub>) of the D form of the microsomal delta-12 desaturase structural gene. Cells (e.g., seeds) of this line can be mutagenized to induce a mutation in a delta-15 desaturase gene, resulting in a plant or plant line carrying a mutation in a delta-12 fatty acid desaturase gene and a mutation in a delta-15 fatty acid desaturase gene.

15 Progeny includes descendants of a particular plant or plant line, e.g., seeds developed on an instant plant are descendants. Progeny of an instant plant include seeds formed on F<sub>1</sub>, F<sub>2</sub>, F<sub>3</sub>, and subsequent generation plants, or seeds formed on BC<sub>1</sub>, BC<sub>2</sub>, BC<sub>3</sub> and subsequent generation plants.

Plants according to the invention preferably contain an altered fatty acid  
20 composition. For example, oil obtained from seeds of such plants may have from about 69 to about 90% oleic acid, based on the total fatty acid composition of the seed. Such oil preferably has from about 74 to about 90% oleic acid, more preferably from about 80 to about 90% oleic acid. In some embodiments, oil obtained from seeds produced by plants of the invention may have from about  
25 2.0% to about 5.0% saturated fatty acids, based on total fatty acid composition of the seeds. In some embodiments, oil obtained from seeds of the invention may have from about 1.0% to about 14.0% linoleic acid, or from about 0.5% to about 10.0%  $\alpha$ -linolenic acid.

Oil composition typically is analyzed by crushing and extracting fatty  
30 acids from bulk seed samples (e.g., 10 seeds). Fatty acid triglycerides in the seed

- 20 -

are hydrolyzed and converted to fatty acid methyl esters. Those seeds having an altered fatty acid composition may be identified by techniques known to the skilled artisan, e.g., gas-liquid chromatography (GLC) analysis of a bulked seed sample, single seed or a single half-seed. Half-seed analysis is well known in the art to be  
5 useful because the viability of the embryo is maintained and thus those seeds having a desired fatty acid profile may be planted to form the next generation. However, half-seed analysis is also known to be an inaccurate representation of genotype of the seed being analyzed. Bulk seed analysis typically yields a more accurate representation of the fatty acid profile of a given genotype. Half-seed  
10 analysis of a population of seeds is, however, a reliable indicator of the likelihood of obtaining a desired fatty acid profile. Fatty acid composition can also be determined on larger samples, e.g., oil obtained by pilot plant or commercial scale refining, bleaching and deodorizing of endogenous oil in the seeds.

The nucleic acid fragments of the invention can be used as markers in  
15 plant genetic mapping and plant breeding programs. Such markers may include restriction fragment length polymorphism (RFLP), random amplification polymorphism detection (RAPD), polymerase chain reaction (PCR) or self-sustained sequence replication (3SR) markers, for example. Marker-assisted breeding techniques may be used to identify and follow a desired fatty acid  
20 composition during the breeding process. Marker-assisted breeding techniques may be used in addition to, or as an alternative to, other sorts of identification techniques. An example of marker-assisted breeding is the use of PCR primers that specifically amplify a sequence containing a desired mutation in delta-12 desaturase or delta-15 desaturase.

25 Methods according to the invention are useful in that the resulting plants and plant lines have desirable seed fatty acid compositions as well as superior agronomic properties compared to known lines having altered seed fatty acid composition. Superior agronomic characteristics include, for example, increased seed germination percentage, increased seedling vigor, increased resistance to

- 21 -

seedling fungal diseases (damping off, root rot and the like), increased yield, and improved standability.

In another aspect, *Brassica* plants producing seeds having a long chain monounsaturated fatty acid content of at least about 82% and an erucic acid content  
5 of at least about 15%, based on total fatty acid composition, are featured. As used herein, "long chain" refers to carbon chains of 16 and greater, e.g., chains of 16 to 24 carbons. The long chain monounsaturated fatty acid content is distributed primarily among oleic acid, eicosenoic acid and erucic acid. The heterogenous nature of the long chain monounsaturated fatty acids in the seed oil triacylglycerols  
10 confers desirable properties to the oil. The levels of total saturated fatty acids and/or the total polyunsaturated fatty acids can be decreased in order to increase the long chain monounsaturate content, i.e., oleic acid, eicosenoic acid, erucic acid and nervonic acid.

High oleic acid lines described herein can be crossed to high erucic acid  
15 lines to produce *Brassica* plants having a high long chain monounsaturated fatty acid content within their seeds. Suitable high oleic acid lines are described, for example, in Example 5 and Table 17, and have an oleic acid content of about 82% to about 85%, based on total fatty acid composition. Suitable high erucic acid lines have an erucic acid content of about 45% or greater, based on total fatty acid  
20 composition. *Brassica* plant line HEC01 is a high erucic acid line that is particularly useful and is sold under the trade name Hero. Other high erucic acid varieties are also known, such as varieties designated Venus, Mercury, Neptune, S89-3673, Dwarf Essex, Reston, Bridger or R-500. McVetty, P.B.E. et al., Can. J. Plant Sci., 76(2):341-342 (1996); Scarth, R. et al., Can. J. Plant Sci., 75(1):205-206  
25 (1995); and McVetty, P.B.E. et al., Can. J. Plant Sci., 76(2):343-344 (1996).

Seeds of the invention can have an oleic acid and eicosenoic acid content of at least about 37% and 14%, respectively, based on total fatty acid composition. The total saturated fatty acid content is less than about 7%. As used herein, "total saturated fatty acid content" refers to the total of myristate (14:0),  
30 palmitate (16:0), stearate (18:0), arachidate (20:0), behenate (22:0) and lignocerate

- 22 -

(24:0). The total polyunsaturated content is less than about 11% based on total fatty acid composition. As used herein, "total polyunsaturated fatty acid content" refers to the sum of linoleic (18:2),  $\alpha$ -linolenic (18:3), and eicosadienoic (20:2) fatty acids as a percentage of the total fatty acid content.

5           In some embodiments, the long chain monounsaturate content in seeds is from about 85% to about 90%. The oleic acid content within these seeds is about 42% or greater, and preferably from about 47% to about 56%. The erucic acid and eicosenoic acid content are from about 17% to about 31% and from about 15% to about 21%, respectively.

10           In some embodiments, the long chain monounsaturate content in seeds is from about 82% to about 90%. The oleic acid content within these seeds is about 25% to about 33%. The erucic acid and eicosenoic acid content are from about 44% to about 50% and from about 10% to about 13%, respectively.

Seed oils having a long chain monounsaturated content of at least about  
15 82% and an erucic acid content of at least about 15%, based on total fatty acid composition, are also featured. These oils can be extracted, for example, from a single line of *Brassica* seeds having a suitable fatty acid composition as described herein. The oleic acid and eicosenoic acid content of these oils is at least about 37% and 14%, respectively, based on total fatty acid composition. The total  
20 saturated and polyunsaturated content of these oils is less than about 7% and 11%, respectively. Preferably, the polyunsaturated content is less than about 9%. In some embodiments, the oils have a monounsaturated content of from about 85% to about 90%. The oleic acid content of these oils is at least about 42% and more preferably, from about 47% to about 56%. The oils have an erucic acid content of  
25 from about 17% to about 31% and an eicosenoic acid content of from about 15% to about 21%.

In some embodiments, the long chain monounsaturate content in such oils is from about 82% to about 90%, comprising an oleic acid content of about 25% to about 33%, an erucic acid of about 44% to about 50% and an eicosenoic  
30 acid content of about 10% to about 13%. The sum of the nervonic acid, erucic

- 23 -

acid and eicosenoic acid contents in such oils can be from about 50% to about 66%.

Alternatively, it is contemplated that oils of the invention can be obtained by mixing high-erucic acid rapeseed oil (HEAR) and an oil having at least  
5 about 87% oleic acid, preferably from about 90% to about 95% oleic acid, based on total fatty acid composition. HEAR oil has an erucic acid content of about 49% and an oleic acid content of about 16%.

Oils having a long chain monounsaturated content of at least about 82% unexpectedly have low temperature properties that are desirable for industrial  
10 applications such as lubrication. The basis for these properties is not known, but it is possible that the heterogeneous chain lengths of the triacylglycerols in oils of the invention impede orderly packing as the end methyl groups have a mismatch in molecular volume, reducing Van der Waals interactions. The double bond in each fatty acid moiety is present at different carbon positions along the acyl chain,  
15 which may disrupt packing and also reduce  $\pi$ - $\pi$  electronic interactions between adjacent fatty acid chains. The high monounsaturate content is thought to provide improved oxidative stability along with high fluidity characteristics. The low levels of polyunsaturates in oils of the invention also promote high oxidative stability, since the rates of oxidation of linoleic acid and linolenic acid at 20°C are 12-20  
20 times and 25 times, respectively, larger than the rate of oxidation of oleic acid.

Oxidative stability can be measured with an Oxidative Stability Index Instrument, Omnion, Inc., Rockland, MA, according to AOCS Official Method Cd 12b-92 (revised 1993). The method is an automated replacement for the Active Oxygen Method (AOM) procedure, AOCS Official Method Cd 12-57. Oxidative  
25 stability of oils having a long chain monounsaturated content of at least about 82% is from about 40 AOM hours to about 100 AOM hours in the absence of added antioxidants. In comparison, mid-oleic canola oil (about 76% oleic acid) and high erucic acid rapeseed oil have oxidative stabilities of about 38 and 16 AOM hours, respectively, in the absence of added antioxidants.

- 24 -

The oils of the invention have desirable functional properties, e.g., low temperature behavior and a high viscosity index, along with high oxidative stability. The presence of higher molecular weight fatty acids increases the molecular weight of the triacylglycerols, providing the oil with a higher flash point and a higher fire point. The increased molecular weight also improves the viscosity index of the oils. Viscosity index is an arbitrary number that indicates the viscosity change with temperature of a lubricant. The Dean and Davis viscosity index can be calculated from observed viscosities of a lubricant at 40°C and 100°C and can produce values ranging from 0 to values greater than 200. A higher viscosity index value indicates that the viscosity of the oil changes less with a change in temperature. In other words, the higher the viscosity index, the smaller the difference in viscosity between high and low temperatures.

An oil of the invention can be formulated for industrial applications such as engine lubricants or as hydraulic fluids by addition of one or more additives to an oil having a long chain monounsaturated fatty acid content of at least about 82% and an erucic acid content of at least about 15%, based on total fatty acid composition. For example, a transmission fluid for diesel engines can be made that includes antioxidants, anti-foam additives, anti-wear additives, corrosion inhibitors, dispersants, detergents, and acid neutralizers, or combinations thereof. Hydraulic oil compositions can include antioxidants, anti-rust additives, anti-wear additives, pour point depressants, viscosity-index improvers and anti-foam additives or combinations thereof. Specific formulations will vary depending on the end use of the oil; suitability of a formulation for a specific end use can be assessed using standard techniques.

Typical antioxidants include zinc dithiophosphates, methyl dithiocarbamates, hindered phenols, phenol sulfides, metal phenol sulfides, metal salicylates, aromatic amines, phospho-sulfurized fats and olefins, sulfurized olefins, sulfurized fats and fat derivatives, sulfurized paraffins, sulfurized carboxylic acids, disalicylal-1,2,-propane diamine, 2,4-bis (alkyldithio-1,3,4-thiadiazoles) and dialkyl selenide. Antioxidants are typically present in amounts from about 0.01% to about

- 25 -

5%, based on the weight of the composition. In particular, about 0.01% to about 1.0% of antioxidant is added to an oil of the invention. See U.S. Patent No. 5,451,334 for additional antioxidants.

Rust inhibitors protect surfaces against rust and include, for example, alkylsuccinic type organic acids, and derivatives thereof, alkylthioacetic acids and derivatives thereof, organic amines, organic phosphates, polyhydric alcohols and sodium and calcium sulphonates. Anti-wear additives adsorb on metal and provide a film that reduces metal-to-metal contact. In general, anti-wear additives include zinc dialkyldithiophosphates, tricresyl phosphate, didodecyl phosphite, sulfurized sperm oil, sulfurized terpenes and zinc dialkyldithiocarbamate, and are used in amounts from about 0.05% to about 4.5%.

Corrosion inhibitors include dithiophosphates and in particular, zinc dithiophosphates, metal sulfonates, metal phenate sulfides, fatty acids, acid phosphate esters and alkyl succinic acids.

Pour point depressants permit flow of the oil composition below the pour point of the unmodified lubricant. Common pour point depressants include polymethacrylates, wax alkylated naphthalene polymers, wax alkylated phenol polymers and chlorinated polymers and are typically present in amounts of about 1% or less. See, for example, U.S. Patent Nos. 5,451,334 and 5,413,725. The viscosity-index can be increased by adding polyisobutylenes, polymethacrylates, polyacrylates, ethylene propylene copolymers, styrene isoprene copolymers, styrene butadiene copolymers and styrene maleic ester copolymers.

Anti-foam additives reduce or prevent the formation of a stable surface foam and are typically present in amounts from about 0.00003% to about 0.05%.

Polymethylsiloxanes, polymethacrylates, salts of alkyl alkylene dithiophosphates, amyl acrylate telomer and poly(2-ethylhexylacrylate-co-ethyl acrylate) are non-limiting examples of anti-foam additives.

Detergents and dispersants are polar materials that provide a cleaning function. Detergents include metal sulfonates, metal salicylates and metal thiophosphonates. Dispersants include polyamine succinimides, hydroxy benzyl

- 26 -

polyamines, polyamine succinamides, polyhydroxy succinic esters and polyamine amide imidazolines.

While the invention is susceptible to various modifications and alternative forms, certain specific embodiments thereof are described in the general methods and examples set forth below. For example the invention may be applied to all *Brassica* species, including *B. rapa*, *B. juncea*, and *B. hirta*, to produce substantially similar results. It should be understood, however, that these examples are not intended to limit the invention to the particular forms disclosed but, instead the invention is to cover all modifications, equivalents and alternatives falling within the scope of the invention. This includes the use of somaclonal variation; physical or chemical mutagenesis of plant parts; anther, microspore or ovary culture followed by chromosome doubling; or self- or cross-pollination to transmit the fatty acid trait, alone or in combination with other traits, to develop new *Brassica* lines.

#### EXAMPLE 1

##### Mutagenesis

Seeds of Westar, a Canadian (*Brassica napus*) spring canola variety, were subjected to chemical mutagenesis. Westar is a registered Canadian spring variety with canola quality. The fatty acid composition of field-grown Westar, 3.9% C<sub>16:0</sub>, 1.9% C<sub>18:0</sub>, 67.5% C<sub>18:1</sub>, 17.6% C<sub>18:2</sub>, 7.4% C<sub>18:3</sub>, <2% C<sub>20:1</sub> + C<sub>22:1</sub>, has remained stable under commercial production, with <± 10% deviation, since 1982.

Prior to mutagenesis, 30,000 seeds of *B. napus* cv. Westar seeds were preimbibed in 300-seed lots for two hours on wet filter paper to soften the seed coat. The preimbibed seeds were placed in 80 mM ethylmethanesulfonate (EMS) for four hours. Following mutagenesis, the seeds were rinsed three times in distilled water. The seeds were sown in 48-well flats containing Pro-Mix. Sixty-eight percent of the mutagenized seed germinated. The plants were maintained at 25°C/15°C, 14/10 hr day/night conditions in the greenhouse. At flowering, each plant was individually self-pollinated.



- 27 -

$M_2$  seed from individual plants were individually catalogued and stored, approximately 15,000  $M_2$  lines was planted in a summer nursery in Carman, Manitoba. The seed from each selfed plant were planted in 3-meter rows with 6-inch row spacing. Westar was planted as the check variety. Selected lines in the field were selfed by bagging the main raceme of each plant. At maturity, the selfed plants were individually harvested and seeds were catalogued and stored to ensure that the source of the seed was known.

Self-pollinated  $M_3$  seed and Westar controls were analyzed in 10-seed bulk samples for fatty acid composition via gas chromatography. Statistical thresholds for each fatty acid component were established using a Z-distribution with a stringency level of 1 in 10,000. Mean and standard deviation values were determined from the non-mutagenized Westar control population in the field. The upper and lower statistical thresholds for each fatty acid were determined from the mean value of the population  $\pm$  the standard deviation, multiplied by the Z-distribution. Based on a population size of 10,000, the confidence interval is 99.99%.

The selected  $M_3$  seeds were planted in the greenhouse along with Westar controls. The seed was sown in 4-inch pots containing Pro-Mix soil and the plants were maintained at 25°C/15°C, 14/10 hr day/night cycle in the greenhouse. At flowering, the terminal raceme was self-pollinated by bagging. At maturity, selfed  $M_4$  seed was individually harvested from each plant, labelled, and stored to ensure that the source of the seed was known.

The  $M_4$  seed was analyzed in 10-seed bulk samples. Statistical thresholds for each fatty acid component were established from 259 control samples using a Z-distribution of 1 in 800. Selected  $M_4$  lines were planted in a field trial in Carman, Manitoba in 3-meter rows with 6-inch spacing. Ten  $M_4$  plants in each row were bagged for self-pollination. At maturity, the selfed plants were individually harvested and the open pollinated plants in the row were bulk harvested. The  $M_5$  seed from single plant selections was analyzed in 10-seed bulk samples and the bulk row harvest in 50-seed bulk samples.

- 28 -

Selected  $M_5$  lines were planted in the greenhouse along with Westar controls. The seed was grown as previously described. At flowering the terminal raceme was self-pollinated by bagging. At maturity, selfed  $M_6$  seed was individually harvested from each plant and analyzed in 10-seed bulk samples for

5 fatty acid composition.

Selected  $M_6$  lines were entered into field trials in Eastern Idaho. The four trial locations were selected for the wide variability in growing conditions. The locations included Burley, Tetonia, Lamont and Shelley (Table 7). The lines were planted in four 3-meter rows with an 8-inch spacing, each plot was replicated

10 four times. The planting design was determined using a Randomized Complete Block Design. The commercial cultivar Westar was used as a check cultivar. At maturity the plots were harvested to determine yield. Yield of the entries in the trial was determined by taking the statistical average of the four replications. The Least Significant Difference Test was used to rank the entries in the randomized

15 complete block design.

TABLE 7Trial Locations for Selected Fatty Acid Mutants

<u>LOCATION</u>	<u>SITE CHARACTERIZATIONS</u>
BURLEY	Irrigated. Long season. High temperatures during flowering.
20 TETONIA	Dryland. Short season. Cool temperatures.
LAMONT	Dryland. Short season. Cool temperatures.
SHELLEY	Irrigated. Medium season. High temperatures during flowering.

To determine the fatty acid profile of entries, plants in each plot were bagged for self-pollination. The  $M_7$  seed from single plants was analyzed for fatty

25 acids in ten-seed bulk samples.

- 29 -

To determine the genetic relationships of the selected fatty acid mutants crosses were made. Flowers of  $M_6$  or later generation mutations were used in crossing.  $F_1$  seed was harvested and analyzed for fatty acid composition to determine the mode of gene action. The  $F_1$  progeny were planted in the  
5 greenhouse. The resulting plants were self-pollinated, the  $F_2$  seed harvested and analyzed for fatty acid composition for allelism studies. The  $F_2$  seed and parent line seed was planted in the greenhouse, individual plants were self-pollinated. The  $F_3$  seed of individual plants was tested for fatty acid composition using 10-seed bulk samples as described previously.

10 In the analysis of some genetic relationships dihaploid populations were made from the microspores of the  $F_1$  hybrids. Self-pollinated seed from dihaploid plants were analyzed for fatty acid analysis using methods described previously.

For chemical analysis, 10-seed bulk samples were hand ground with a glass rod in a 15-mL polypropylene tube and extracted in 1.2 mL 0.25 N KOH in  
15 1:1 ether/methanol. The sample was vortexed for 30 sec. and heated for 60 sec. in a 60°C water bath. Four mL of saturated NaCl and 2.4 mL of iso-octane were added, and the mixture was vortexed again. After phase separation, 600  $\mu$ L of the upper organic phase were pipetted into individual vials and stored under nitrogen at -5°C. One  $\mu$ L samples were injected into a Supelco SP-2330 fused silica capillary  
20 column (0.25 mm ID, 30 M length, 0.20  $\mu$ m df).

The gas chromatograph was set at 180°C for 5.5 minutes, then programmed for a 2°C/minute increase to 212°C, and held at this temperature for 1.5 minutes. Total run time was 23 minutes. Chromatography settings were:  
Column head pressure - 15 psi, Column flow (He) - 0.7 mL/min., Auxiliary and  
25 Column flow - 33 mL/min., Hydrogen flow - 33 mL/min., Air flow - 400 mL/min., Injector temperature - 250°C, Detector temperature - 300°C, Split vent - 1/15.

Table 8 describes the upper and lower statistical thresholds for each fatty acid of interest.

- 30 -

TABLE 8

Statistical Thresholds for Specific Fatty Acids  
Derived from Control Westar Plantings

		Percent Fatty Acids					
5	Genotype	C <sub>16:0</sub>	C <sub>18:0</sub>	C <sub>18:1</sub>	C <sub>18:2</sub>	C <sub>18:3</sub>	Sats*
	<hr/>						
	M <sub>3</sub> Generation(1 in 10,000 rejection rate)						
	Lower	3.3	1.4	--	13.2	5.3	6.0
	Upper	4.3	2.5	71.0	21.6	9.9	8.3
10	<hr/>						
	M <sub>4</sub> Generation(1 in 800 rejection rate)						
	Lower	3.6	0.8	--	12.2	3.2	5.3
	Upper	6.3	3.1	76.0	32.4	9.9	11.2
	<hr/>						
	M <sub>5</sub> Generation (1 in 755 rejection rate)						
	Lower	2.7	0.9	--	9.6	2.6	4.5
15	Upper	5.7	2.7	80.3	26.7	9.6	10.0

\*Sats=Total Saturate Content

EXAMPLE 2High Oleic Acid Canola Lines

In the studies of Example 1, at the M<sub>3</sub> generation, 31 lines exceeded the upper statistical threshold for oleic acid ( $\geq 71.0\%$ ). Line W7608.3 had 71.2% oleic acid. At the M<sub>4</sub> generation, its selfed progeny (W7608.3.5, since designated A129.5) continued to exceed the upper statistical threshold for C<sub>18:1</sub> with 78.8% oleic acid. M<sub>5</sub> seed of five self-pollinated plants of line A129.5 (ATCC 40811) averaged 75.0% oleic acid. A single plant selection, A129.5.3 had 75.6% oleic acid. The fatty acid composition of this high oleic acid mutant, which was stable under both field and greenhouse conditions to the M<sub>7</sub> generation, is summarized in Table 9. This line also stably maintained its mutant fatty acid composition to the M<sub>7</sub> generation in field trials in multiple locations. Over all locations the self-pollinated plants (A129) averaged 78.3% oleic acid. The fatty acid composition of the A129 for each Idaho trial location are summarized in Table 10. In multiple location replicated yield trials, A129 was not significantly different in yield from the parent cultivar Westar.

- 31 -

The canola oil of A129, after commercial processing, was found to have superior oxidative stability compared to Westar when measured by the Accelerated Oxygen Method (AOM), American Oil Chemists' Society Official Method Cd 12-57 for fat stability; Active Oxygen Method (revised 1989). The AOM of Westar was 18 AOM hours and for A129 was 30 AOM hours.

**TABLE 9**

Fatty Acid Composition of a High  
Oleic Acid Canola Line Produced by Seed Mutagenesis

		Percent Fatty Acids					
10	Genotype	C <sub>16:0</sub>	C <sub>18:0</sub>	C <sub>18:1</sub>	C <sub>18:2</sub>	C <sub>18:3</sub>	Sats
	Westar	3.9	1.9	67.5	17.6	7.4	7.0
	W7608.3 (M <sub>3</sub> )	3.9	2.4	71.2	12.7	6.1	7.6
	W7608.3.5 (M <sub>4</sub> )	3.9	2.0	78.8	7.7	3.9	7.3
15	A129.5.3 (M <sub>5</sub> )	3.8	2.3	75.6	9.5	4.9	7.6

Sats=Total Saturate Content

**TABLE 10**

Fatty Acid Composition of a Mutant High  
Oleic Acid Line at Different Field Locations in Idaho

		Percent Fatty Acids					
Location		C <sub>16:0</sub>	C <sub>18:0</sub>	C <sub>18:1</sub>	C <sub>18:2</sub>	C <sub>18:3</sub>	Sats
25	Burley	3.3	2.1	77.5	8.1	6.0	6.5
	Tetonia	3.5	3.4	77.8	6.5	4.7	8.5
	Lamont	3.4	1.9	77.8	7.4	6.5	6.3
	Shelley	3.3	2.6	80.0	5.7	4.5	7.7

Sats=Total Saturate Content

- 32 -

The genetic relationship of the high oleic acid mutation A129 to other oleic desaturases was demonstrated in crosses made to commercial canola cultivars and a low linolenic acid mutation. A129 was crossed to the commercial cultivar Global ( $C_{16:0}$  - 4.5%,  $C_{18:0}$  - 1.5%,  $C_{18:1}$  - 62.9%,  $C_{18:2}$  - 20.0%,  $C_{18:3}$  - 7.3%).

- 5 Approximately 200  $F_2$  individuals were analyzed for fatty acid composition. The results are summarized in Table 11. The segregation fit 1:2:1 ratio suggesting a single co-dominant gene controlled the inheritance of the high oleic acid phenotype.

TABLE 11

Genetic Studies of A129 X Global

Frequency

10

Genotype	$C_{18:1}$	Observed	Expected
	Content(%)		
od-od-	77.3	43	47
od-od+	71.7	106	94
15 od+od+	66.1	49	47

- A cross between A129 and IMC 01, a low linolenic acid variety ( $C_{16:0}$  - 4.1%,  $C_{18:0}$  - 1.9%,  $C_{18:1}$  - 66.4%,  $C_{18:2}$  - 18.1%,  $C_{18:3}$  - 5.7%), was made to determine the inheritance of the oleic acid desaturase and linoleic acid desaturase. In the  $F_1$  hybrids both the oleic acid and linoleic acid desaturase genes approached the mid-parent values indicating a co-dominant gene actions. Fatty acid analysis of the  $F_2$  individuals confirmed a 1:2:1:2:4:2:1:2:1 segregation of two independent, co-dominant genes (Table 12). A line was selected from the cross of A129 and IMC01 and designated as IMC130 (ATCC deposit no. 75446) as described in U.S. Patent Application No. 08/425,108, incorporated herein by reference.

- 33 -

**TABLE 12**Genetic Studies of A129 X IMC 01

		<u>Frequency</u>	
<u>Genotype</u>	<u>Ratio</u>	<u>Observed</u>	<u>Expected</u>
5 od-od-ld-ld-	1	11	12
od-od-ld-ld+	2	30	24
od-od-ld+ld+	1	10	12
od-od+ld-ld-	2	25	24
od-od+ld-ld+	4	54	47
10 od-od+ld+ld+	2	18	24
od+od+ld-ld-	1	7	12
od+od+ld-ld+	2	25	24
od+od+ld+ld+	1	8	12

An additional high oleic acid line, designated A128.3, was also produced  
 15 by the disclosed method. A 50-seed bulk analysis of this line showed the following  
 fatty acid composition: C<sub>16:0</sub> - 3.5%, C<sub>18:0</sub> - 1.8%, C<sub>18:1</sub> - 77.3%, C<sub>18:2</sub> - 9.0%, C<sub>18:3</sub> -  
 5.6%, FDA Sats - 5.3%, Total Sats - 6.4%. This line also stably maintained its  
 mutant fatty acid composition to the M<sub>7</sub> generation. In multiple locations  
 replicated yield trials, A128 was not significantly different in yield from the parent  
 20 cultivar Westar.

A129 was crossed to A128.3 for allelism studies. Fatty acid  
 composition of the F<sub>2</sub> seed showed the two lines to be allelic. The mutational  
 events in A129 and A128.3 although different in origin were in the same gene.

An additional high oleic acid line, designated M3028.-10 (ATCC  
 25 75026), was also produced by the disclosed method in Example 1. A 10-seed bulk  
 analysis of this line showed the following fatty acid composition: C<sub>16:0</sub> - 3.5%, C<sub>18:0</sub>  
 - 1.8%, C<sub>18:1</sub> - 77.3%, C<sub>18:2</sub> - 9.0%, C<sub>18:3</sub> - 5.6%, FDA Saturates - 5.3%, Total  
 Saturates - 6.4%. In a single location replicated yield trial M3028.10 was not  
 significantly different in yield from the parent cultivar Westar.

- 34 -

EXAMPLE 3Low Linoleic Acid Canola

In the studies of Example 1, at the  $M_3$  generation, 80 lines exceeded the lower statistical threshold for linoleic acid ( $\leq 13.2\%$ ). Line W12638.8 had 9.4% linoleic acid. At the  $M_4$  and  $M_5$  generations, its selfed progenies [W12638.8, since designated A133.1 (ATCC 40812)] continued to exceed the statistical threshold for low  $C_{18:2}$  with linoleic acid levels of 10.2% and 8.4%, respectively. The fatty acid composition of this low linoleic acid mutant, which was stable to the  $M_7$  generation under both field and greenhouse conditions, is summarized in Table 13. In multiple location replicated yield trials, A133 was not significantly different in yield from the parent cultivar Westar. An additional low linoleic acid line, designated M3062.8 (ATCC 75025), was also produced by the disclosed method. A 10-seed bulk analysis of this line showed the following fatty acid composition:  $C_{16:0}$  - 3.8%,  $C_{18:0}$  - 2.3%,  $C_{18:1}$  - 77.1%,  $C_{18:2}$  - 8.9%,  $C_{18:3}$  - 4.3%, FDA Sats-6.1%. This line has also stably maintained its mutant fatty acid composition in the field and greenhouse.

TABLE 13

Fatty Acid Composition of a Low  
Linoleic Acid Canola Line Produced by Seed Mutagenesis

20	Percent Fatty Acids						
	Genotype	C <sub>16:0</sub>	C <sub>18:0</sub>	C <sub>18:1</sub>	C <sub>18:2</sub>	C <sub>18:3</sub>	Sats <sup>b</sup>
	Westar	3.9	1.9	67.5	17.6	7.4	7.0
	W12638.8 (M <sub>3</sub> )	3.9	2.3	75.0	9.4	6.1	7.5
25	W12638.8.1 (M <sub>4</sub> )	4.1	1.7	74.6	10.2	5.9	7.1
	A133.1.8 (M <sub>5</sub> )	3.8	2.0	77.7	8.4	5.0	7.0

<sup>a</sup>Letter and numbers up to second decimal point indicate the plant line. Number after second decimal point indicates an individual plant.

<sup>b</sup>Sats=Total Saturated Content



- 35 -

EXAMPLE 4Low Linolenic and Linoleic Acid Canola

In the studies of Example 1, at the M<sub>3</sub> generation, 57 lines exceeded the lower statistical threshold for linolenic acid ( $\leq 5.3\%$ ). Line W14749.8 had 5.3% linolenic acid and 15.0% linoleic acid. At the M<sub>4</sub> and M<sub>5</sub> generations, its selfed progenies [W14749.8, since designated M3032 (ATCC 75021)] continued to exceed the statistical threshold for low C<sub>18:3</sub> with linolenic acid levels of 2.7% and 2.3%, respectively, and for a low sum of linolenic and linoleic acids with totals of 11.8% and 12.5% respectively. The fatty acid composition of this low linolenic acid plus linoleic acid mutant, which was stable to the M<sub>5</sub> generation under both field and greenhouse conditions, is summarized in Table 14. In a single location replicated yield trial M3032 was not significantly different in yield from the parent cultivar (Westar).

TABLE 14

Fatty Acid Composition of a Low  
Linolenic Acid Canola Line Produced by Seed Mutagenesis  
Percent Fatty Acids

Genotype	C <sub>16:0</sub>	C <sub>18:0</sub>	C <sub>18:1</sub>	C <sub>18:2</sub>	C <sub>18:3</sub>	Sats
Westar	3.9	1.9	67.5	17.6	7.4	7.0
W14749.8 (M <sub>3</sub> )	4.0	2.5	69.4	15.0	5.3	6.5
M3032.8 (M <sub>4</sub> )	3.9	2.4	77.9	9.1	2.7	6.4
M3032.1 (M <sub>5</sub> )	3.5	2.8	80.0	10.2	2.3	6.5

Sats=Total Saturate Content

EXAMPLE 5Canola Lines Q508 and Q4275

Seeds of the *B. napus* line IMC-129 were mutagenized with methyl N-nitrosoguanidine (MNNG). The MNNG treatment consisted of three parts: pre-

- 36 -

soak, mutagen application, and wash. A 0.05M Sorenson's phosphate buffer was used to maintain pre-soak and mutagen treatment pH at 6.1. Two hundred seeds were treated at one time on filter paper (Whatman #3M) in a petri dish (100mm x 15mm). The seeds were pre-soaked in 15 mls of 0.05M Sorenson's buffer, pH 6.1, 5 under continued agitation for two hours. At the end of the pre-soak period, the buffer was removed from the plate.

A 10mM concentration of MNNG in 0.05M Sorenson's buffer, pH 6.1, was prepared prior to use. Fifteen ml of 10m MNNG was added to the seeds in each plate. The seeds were incubated at  $22^{\circ}\text{C} \pm 3^{\circ}\text{C}$  in the dark under constant 10 agitation for four (4) hours. At the end of the incubation period, the mutagen solution was removed.

The seeds were washed with three changes of distilled water at 10 minute intervals. The fourth wash was for thirty minutes. This treatment regime produced an LD60 population.

15 Treated seeds were planted in standard greenhouse potting soil and placed into an environmentally controlled greenhouse. The plants were grown under sixteen hours of light. At flowering, the racemes were bagged to produce selfed seed. At maturity, the M2 seed was harvested. Each M2 line was given an identifying number. The entire MNNG-treated seed population was designated as 20 the Q series.

Harvested M2 seeds was planted in the greenhouse. The growth conditions were maintained as previously described. The racemes were bagged at flowering for selfing. At maturity, the selfed M3 seed was harvested and analyzed for fatty acid composition. For each M3 seed line, approximately 10-15 seeds were 25 analyzed in bulk as described in Example 1.

High oleic-low linoleic M3 lines were selected from the M3 population using a cutoff of >82% oleic acid and <5.0% linoleic. From the first 1600 M3 lines screened for fatty acid composition, Q508 was identified. The Q508 M3 generation was advanced to the M4 generation in the greenhouse. Table 15 shows

- 37 -

the fatty acid composition of Q508 and IMC 129. The M4 selfed seed maintained the selected high oleic-low linoleic acid phenotype (Table 16).

**TABLE 15**

Fatty Acid Composition of A129 and High  
Oleic Acid M3 Mutant Q508

Line #	16:0	18:0	18:1	18:2	18:3
A129*	4.0	2.4	77.7	7.8	4.2
Q508	3.9	2.1	84.9	2.4	2.9

\*Fatty acid composition of A129 is the average of  
50 self-pollinated plants grown with the M3 population

M<sub>4</sub> generation Q508 plants had poor agronomic qualities in the field compared to Westar. Typical plants were slow growing relative to Westar, lacked early vegetative vigor, were short in stature, tended to be chlorotic and had short pods. The yield of Q508 was very low compared to Westar.

The M<sub>4</sub> generation Q508 plants in the greenhouse tended to be reduced in vigor compared to Westar. However, Q508 yields in the greenhouse were greater than Q508 yields in the field.

**TABLE 16**

Fatty Acid Composition of Seed Oil  
from Greenhouse-Grown Q508, IMC 129 and Westar.

Line	16:0	18:0	18:1	18:2	18:3	FDA Sats
IMC 129 <sup>a</sup>	4.0	2.4	77.7	7.8	4.2	6.4
Westar <sup>b</sup>	3.9	1.9	67.5	17.6	7.4	>5.8
Q508 <sup>c</sup>	3.9	2.1	84.9	2.4	2.9	6.0

<sup>a</sup>Average of 50 self-pollinated plants

<sup>b</sup>Data from Example 1

<sup>c</sup>Average of 50 self-pollinated plants

- 38 -

Nine other M<sub>4</sub> high-oleic low-linoleic lines were also identified: Q3603, Q3733, Q4249, Q6284, Q6601, Q6761, Q7415, Q4275, and Q6676. Some of these lines had good agronomic characteristics and an elevated oleic acid level in seeds of about 80% to about 84%.

- 5 Q4275 was crossed to the variety Cyclone. After selfing for seven generations, mature seed was harvested from 93GS34-179, a progeny line of the Q4275x<sup>5</sup>Cyclone cross. Referring to Table 17, fatty acid composition of a bulk seed sample shows that 93GS34 retained the seed fatty acid composition of Q4275. 93GS34-179 also maintained agronomically desirable characteristics.
- 10 After more than seven generations of selfing of Q4275, plants of Q4275, IMC 129 and 93GS34 were field grown during the summer season. The selections were tested in 4 replicated plots (5 feet X 20 feet) in a randomized block design. Plants were open pollinated. No selfed seed was produced. Each plot was harvested at maturity, and a sample of the bulk harvested seed from each line was
- 15 analyzed for fatty acid composition as described above. The fatty acid compositions of the selected lines are shown in Table 17.

TABLE 17

Fatty Acid Composition of  
Field Grown IMC 129, Q4275 and 93GS34 Seeds

20

Line	Fatty Acid Composition (%)					
	C <sub>16:0</sub>	C <sub>18:0</sub>	C <sub>18:1</sub>	C <sub>18:2</sub>	C <sub>18:3</sub>	FDA Sats
IMC 129	3.3	2.4	76.7	8.7	5.2	5.7
Q4275	3.7	3.1	82.1	4.0	3.5	6.8
93GS34-179	2.6	2.7	85.0	2.8	3.3	5.3

- The results shown in Table 17 show that Q4275 maintained the selected
- 25 high oleic - low linoleic acid phenotype under field conditions. The agronomic characteristics of Q4275 plants were superior to those of Q508.

M<sub>4</sub> generation Q508 plants were crossed to a dihaploid selection of Westar, with Westar serving as the female parent. The resulting F<sub>1</sub> seed was

- 39 -

termed the 92EF population. About 126 F<sub>1</sub> individuals that appeared to have better agronomic characteristics than the Q508 parent were selected for selfing. A portion of the F<sub>2</sub> seed from such individuals was replanted in the field. Each F<sub>2</sub> plant was selfed and a portion of the resulting F<sub>3</sub> seed was analyzed for fatty acid composition. The content of oleic acid in F<sub>3</sub> seed ranged from 59 to 79%. No high oleic (>80%) individuals were recovered with good agronomic type.

A portion of the F<sub>2</sub> seed of the 92EF population was planted in the greenhouse to analyze the genetics of the Q508 line. F<sub>3</sub> seed was analyzed from 380 F<sub>2</sub> individuals. The C<sub>18:1</sub> levels of F<sub>3</sub> seed from the greenhouse experiment is depicted in Figure 1. The data were tested against the hypothesis that Q508 contains two mutant genes that are semi-dominant and additive: the original IMC 129 mutation as well as one additional mutation. The hypothesis also assumes that homozygous Q508 has greater than 85% oleic acid and homozygous Westar has 62-67% oleic acid. The possible genotypes at each gene in a cross of Q508 by Westar may be designated as:

AA = Westar Fad2<sup>a</sup>

BB = Westar Fad2<sup>b</sup>

aa = Q508 Fad2<sup>a</sup>

bb = Q508 Fad2<sup>b</sup>

Assuming independent segregation, a 1:4:6:4:1 ratio of phenotypes is expected. The phenotypes of heterozygous plants are assumed to be indistinguishable and, thus, the data were tested for fit to a 1:14:1 ratio of homozygous Westar: heterozygous plants: homozygous Q508.

Phenotypic	# of	
Ratio	Westar Alleles	Genotype
1	4	AABB(Westar)
4	3	AABb,AaBB,AABb,AaBB
6	2	AaBb,AAbb,AaBb,AaBb,aaBB,AaBb
4	1	Aabb,aaBb,Aabb,aaBb
1	0	aabb (Q508)

- 40 -

Using Chi-square analysis, the oleic acid data fit a 1:14:1 ratio. It was concluded that Q508 differs from Westar by two major genes that are semi-dominant and additive and that segregate independently. By comparison, the genotype of IMC 129 is aaBB.

- 5        The fatty acid composition of representative F3 individuals having greater than 85% oleic acid in seed oil is shown in Table 18. The levels of saturated fatty acids are seen to be decreased in such plants, compared to Westar.

**TABLE 18**  
92EF F<sub>3</sub> Individuals with >85% C<sub>18:1</sub> in Seed Oil

F3 Plant Identifier	Fatty Acid Composition (%)					
	C16:0	C18:0	C18:1	C18:2	C18:3	FDASA
+38068	3.401	1.582	85.452	2.134	3.615	4.983
+38156	3.388	1.379	85.434	2.143	3.701	4.767
+38171	3.588	1.511	85.289	2.367	3.425	5.099
+38181	3.75	1.16	85.312	2.968	3.819	4.977
+38182	3.529	0.985	85.905	2.614	3.926	4.56
+38191	3.364	1.039	85.737	2.869	4.039	4.459
+38196	3.557	1.182	85.054	2.962	4.252	4.739
+38202	3.554	1.105	86.091	2.651	3.721	4.713
+38220	3.093	1.16	86.421	1.931	3.514	4.314
+38236	3.308	1.349	85.425	2.37	3.605	4.718
+38408	3.617	1.607	85.34	2.33	3.562	5.224
+38427	3.494	1.454	85.924	2.206	3.289	4.948
+38533	3.64	1.319	85.962	2.715	3.516	4.959

- 41 -

EXAMPLE 6Leaf and Root Fatty Acid Profiles of Canola  
Lines IMC-129, Q508, and Westar

Plants of Q508, IMC 129 and Westar were grown in the greenhouse.

- 5 Mature leaves, primary expanding leaves, petioles and roots were harvested at the 6-8 leaf stage, frozen in liquid nitrogen and stored at -70°C. Lipid extracts were analyzed by GLC as described in Example 1. The fatty acid profile data are shown in Table 19. The data in Table 19 indicate that total leaf lipids in Q508 are higher in  $C_{18:1}$  content than the  $C_{18:2}$  plus  $C_{18:3}$  content. The reverse is true for Westar and
- 10 IMC 129. The difference in total leaf lipids between Q508 and IMC 129 is consistent with the hypothesis that a second Fad2 gene is mutated in Q508.

- The  $C_{16:3}$  content in the total lipid fraction was about the same for all three lines, suggesting that the plastid FadC gene product was not affected by the Q508 mutations. To confirm that the FadC gene was not mutated, chloroplast
- 15 lipids were separated and analyzed. No changes in chloroplast  $C_{16:1}$ ,  $C_{16:2}$  or  $C_{16:3}$  fatty acids were detected in the three lines. The similarity in plastid leaf lipids among Q508, Westar and IMC 129 is consistent with the hypothesis that the second mutation in Q508 affects a microsomal Fad2 gene and not a plastid FadC gene.

**TABLE 19**

	MATURE LEAF			EXPANDING LEAF			PETIOLE			ROOT		
	West.	129	3Q508	West.	129	3Q508	West.	129	3Q508	West.	129	3Q508
16:0	12.1	11.9	10.1	16.4	16.1	11.3	21.7	23.5	11.9	21.1	21.9	12.0
16:1	0.8	0.6	1.1		0.6	1.1	1.0	1.3	1.4	-	-	-
16:2	2.3	2.2	2.0	2.8	3.1	2.8	1.8	2.2	1.8	-	-	-
16:3	14.7	15.0	14.0	6.3	5.4	6.9	5.7	4.6	5.7	-	-	-
18:0	2.2	1.6	1.2	2.5	2.8	1.5	3.7	4.0	1.6	3.6	2.9	2.5
18:1	2.8	4.9	16.7	3.8	8.3	38.0	4.9	12.9	46.9	3.5	6.1	68.8
18:2	12.6	11.5	6.8	13.3	13.8	4.9	20.7	18.3	5.2	28.0	30.4	4.4
18:3	50.6	50.3	46.0	54.2	50.0	33.5	40.4	33.2	25.3	43.8	38.7	12.3



- 43 -

EXAMPLE 7

Sequences of Mutant and Wild-Type Delta-12 Fatty Acid  
Desaturases from *B. napus*

Primers specific for the FAD2 structural gene were used to clone the  
5 entire open reading frame (ORF) of the D and F delta-12 desaturase genes by  
reverse transcriptase polymerase chain reaction (RT-PCR). RNA from seeds of  
IMC 129, Q508 and Westar plants was isolated by standard methods and was used  
as template. The RT-amplified fragments were used for nucleotide sequence  
determination. The DNA sequence of each gene from each line was determined  
10 from both strands by standard dideoxy sequencing methods.

Sequence analysis revealed a G to A transversion at nucleotide 316  
(from the translation initiation codon) of the D gene in both IMC 129 and Q508,  
compared to the sequence of Westar. The transversion changes the codon at this  
position from GAG to AAG and results in a non-conservative substitution of  
15 glutamic acid, an acidic residue, for lysine a basic residue. The presence of the  
same mutation in both lines was expected since the Q508 line was derived from  
IMC 129. The same base change was also detected in Q508 and IMC 129 when  
RNA from leaf tissue was used as template.

The G to A mutation at nucleotide 316 was confirmed by sequencing  
20 several independent clones containing fragments amplified directly from genomic  
DNA of IMC 129 and Westar. These results eliminated the possibility of a rare  
mutation introduced during reverse transcription and PCR in the RT-PCR protocol.  
It was concluded that the IMC 129 mutant is due to a single base transversion at  
nucleotide 316 in the coding region of the D gene of rapeseed microsomal delta 12-  
25 desaturase.

A single base transition from T to A at nucleotide 515 of the F gene  
was detected in Q508 compared to the Westar sequence. The mutation changes the  
codon at this position from CTC to CAC, resulting in the non-conservative  
substitution of a non-polar residue, leucine, for a polar residue, histidine, in the

- 44 -

resulting gene product. No mutations were found in the F gene sequence of IMC 129 compared to the F gene sequence of Westar.

These data support the conclusion that a mutation in a delta-12 desaturase gene sequence results in alterations in the fatty acid profile of plants containing such a mutated gene. Moreover, the data show that when a plant line or species contains two delta-12 desaturase loci, the fatty acid profile of an individual having two mutated loci differs from the fatty acid profile of an individual having one mutated locus.

The mutation in the D gene of IMC 129 and Q508 mapped to a region having a conserved amino acid motif (His-Xaa-Xaa-Xaa-His) found in cloned delta-12 and delta-15 membrane bound-desaturases (Table 20).

**TABLE 20**

Alignment of Amino Acid Sequences  
of Cloned Canola Membrane Bound-Desaturases

Desaturase Gene	Sequence <sup>a</sup>	Position
Canola-fad2-D(mutant)	AHKCGH	109-114
Canola-Fad2-D	AHECGH	109-114
Canola-Fad2-F	AHECGH	109-114
Canola-FadC	<u>GH</u> <b>D</b> <u>CA</u> H	170-175
Canola-fad3 (mutant)	<u>GH</u> KCGH	94-99
Canola-Fad3	<u>GH</u> <b>D</b> CGH	94-99
Canola-FadD	<u>GH</u> <b>D</b> CGH	125-130

(FadD = Plastid delta 15, Fad3 = Microsomal delta-15),  
(FadC = Plastid delta-12, Fad2 = Microsomal delta-12)

<sup>a</sup> One letter amino acid code; conservative substitutions are underlined; non-conservative substitutions are in bold.

- 45 -

EXAMPLE 8

Transcription and Translation of Microsomal Delta-12  
Fatty Acid Desaturases

Transcription in vivo was analyzed by RT-PCR analysis of stage II and  
5 stage III developing seeds and leaf tissue. The primers used to specifically amplify  
delta-12 desaturase F gene RNA from the indicated tissues were sense primer 5'-  
GGATATGATGATGGTGAAAGA-3' and antisense primer 5'-  
TCTTTCACCATCATCATATCC-3'. The primers used to specifically amplify  
delta-12 desaturase D gene RNA from the indicated tissues were sense primer 5'-  
10 GTTATGAAGCAAAGAAGAAAC-3' and antisense primer 5'-  
GTTTCTTCTTTGCTTCATAAC-3'. The results indicated that mRNA of both the  
D and F gene was expressed in seed and leaf tissues of IMC 129, Q508 and wild  
type Westar plants.

*In vitro* transcription and translation analysis showed that a peptide of  
15 about 46 kD was made. This is the expected size of both the D gene product and  
the F gene product, based on sum of the deduced amino acid sequence of each  
gene and the cotranslational addition of a microsomal membrane peptide.

These results rule out the possibility that non-sense or frameshift  
mutations, resulting in a truncated polypeptide gene product, are present in either  
20 the mutant D gene or the mutant F gene. The data, in conjunction with the data of  
Example 7, support the conclusion that the mutations in Q508 and IMC 129 are in  
delta-12 fatty acid desaturase structural genes encoding desaturase enzymes, rather  
than in regulatory genes.

EXAMPLE 9

25 Development of Gene-Specific PCR Markers

Based on the single base change in the mutant D gene of IMC 129  
described in above, two 5' PCR primers were designed. The nucleotide sequence  
of the primers differed only in the base (G for Westar and A for IMC 129) at the  
3' end. The primers allow one to distinguish between mutant fad2-D and wild-type

- 46 -

Fad2-D alleles in a DNA-based PCR assay. Since there is only a single base difference in the 5' PCR primers, the PCR assay is very sensitive to the PCR conditions such as annealing temperature, cycle number, amount, and purity of DNA templates used. Assay conditions have been established that distinguish  
5 between the mutant gene and the wild type gene using genomic DNA from IMC 129 and wild type plants as templates. Conditions may be further optimized by varying PCR parameters, particularly with variable crude DNA samples. A PCR assay distinguishing the single base mutation in IMC 129 from the wild type gene along with fatty acid composition analysis provides a means to simplify segregation  
10 and selection analysis of genetic crosses involving plants having a delta-12 fatty acid desaturase mutation.

#### EXAMPLE 10

##### Transformation with Mutant and Wild Type Fad3 Genes

*B. napus* cultivar Westar was transformed with mutant and wild type  
15 Fad3 genes to demonstrate that the mutant Fad3 gene for canola cytoplasmic linoleic desaturase delta-15 desaturase is nonfunctional. Transformation and regeneration were performed using disarmed *Agrobacterium tumefaciens* essentially following the procedure described in WO 94/11516.

Two disarmed *Agrobacterium* strains were engineered, each containing a  
20 Ti plasmid having the appropriate gene linked to a seed-specific promoter and a corresponding termination sequence. The first plasmid, pIMC110, was prepared by inserting into a disarmed Ti vector the full length wild type Fad3 gene in sense orientation (nucleotides 208 to 1336 of SEQ ID 6 in WO 93/11245), flanked by a napin promoter sequence positioned 5' to the Fad3 gene and a napin termination  
25 sequence positioned 3' to the Fad3 gene. The rapeseed napin promoter is described in EP 0255378.

The second plasmid, pIMC205, was prepared by inserting a mutated Fad3 gene in sense orientation into a disarmed Ti vector. The mutant sequence contained mutations at nucleotides 411 and 413 of the microsomal Fad3 gene

- 47 -

described in WO93/11245, thus changing the sequence for codon 96 from GAC to AAG. The amino acid at codon 96 of the gene product was thereby changed from aspartic acid to lysine. See Table 20. A bean (*Phaseolus vulgaris*) phaseolin (7S seed storage protein) promoter fragment of 495 base pairs was placed 5' to the  
5 mutant Fad3 gene and a phaseolin termination sequence was placed 3' to the mutant Fad3 gene. The phaseolin sequence is described in Doyle et al., (1986) J. Biol. Chem. 261:9228-9238) and Slightom et al., (1983) Proc. Natl. Acad. Sci. USA 80:1897-1901.

The appropriate plasmids were engineered and transferred separately to  
10 *Agrobacterium* strain LBA4404. Each engineered strain was used to infect 5 mm segments of hypocotyl explants from Westar seeds by cocultivation. Infected hypocotyls were transferred to callus medium and, subsequently, to regeneration medium. Once discernable stems formed from the callus, shoots were excised and transferred to elongation medium. The elongated shoots were cut, dipped in  
15 Rootone™, rooted on an agar medium and transplanted to potting soil to obtain fertile T1 plants. T2 seeds were obtained by selfing the resulting T1 plants.

Fatty acid analysis of T2 seeds was carried out as described above. The results are summarized in Table 21. Of the 40 transformants obtained using the pIMC110 plasmid, 17 plants demonstrated wild type fatty acid profiles and 16  
20 demonstrated overexpression. A proportion of the transformants are expected to display an overexpression phenotype when a functioning gene is transformed in sense orientation into plants.

Of the 307 transformed plants having the pIMC205 gene, none exhibited a fatty acid composition indicative of overexpression. This result indicates that the  
25 mutant fad3 gene product is non-functional, since some of the transformants would have exhibited an overexpression phenotype if the gene product were functional.

**TABLE 21**  
Overexpression and Co-suppression Events in  
Westar Populations Transformed with pIMC205 or pIMC110.

Construct	Number of Transformants	$\alpha$ - Linolenic Acid Range(%)	Overexpression Events ( $>10\%$ linolenic)	Co-Suppression Events ( $<4.0\%$ linolenic)	Wild Type Events
pIMC110	40	2.4 - 20.6	16	7	17
pIMC205	307	4.6 - 10.4	0	0	307

- 49 -

Fatty acid compositions of representative transformed plants are presented in Table 22. Lines 652-09 and 663-40 are representative of plants containing pIMC110 and exhibiting an overexpression and a co-suppression phenotype, respectively. Line 205-284 is representative of plants containing

5 pIMC205 and having the mutant fad3 gene.

**TABLE 22**

Fatty Acid Composition of T2 Seed  
From Westar Transformed With pIMC205 or pIMC110.

Line	Fatty Acid Composition (%)				
	C16:0	C18:0	C18:1	C18:2	C18:3
10 652-09 pIMC110 overexpression	4.7	3.3	65.6	8.1	14.8
663-40 pIMC110 co-suppression	4.9	2.1	62.5	23.2	3.6
15 205-284 pIMC205	3.7	1.8	68.8	15.9	6.7

**EXAMPLE 11****Sequences of Wild Type and Mutant Fad2-D and Fad2-F**

High molecular weight genomic DNA was isolated from leaves of

20 Q4275 plants (Example 5). This DNA was used as template for amplification of Fad2-D and Fad2-F genes by polymerase chain reaction (PCR). PCR amplifications were carried out in a total volume of 100  $\mu$ l and contained 0.3  $\mu$ g genomic DNA, 200  $\mu$ M deoxyribonucleoside triphosphates, 3 mM MgSO<sub>4</sub>, 1-2 Units DNA polymerase and 1X Buffer (supplied by the DNA polymerase

25 manufacturer). Cycle conditions were: 1 cycle for 1 min at 95°C, followed by 30 cycles of 1 min at 94°C, 2 min at 55°C and 3 min at 73°C.

The Fad2-D gene was amplified once using Elongase® (Gibco-BRL). PCR primers were: CAUCAUCAUCAUUCTTCTTCGTAGGGTTCATCG and

- 50 -

CUACUACUACUATCATAGAAGAGAAAGGTTTCAG for the 5' and 3' ends of the gene, respectively.

The Fad2-F gene was independently amplified 4 times, twice with Elongase® and twice with Taq polymerase (Boehringer Mannheim). The PCR  
5 primers used were: 5'CAUCAUCAUCAUCATGGGTGCACGTGGAAGAA3' and 5'CUACUACUACUATCTTTCACCATCATCATATCC3' for the 5' and 3' ends of the gene, respectively.

Amplified DNA products were resolved on an agarose gel, purified by JetSorb® and then annealed into pAMP1 (Gibco-BRL) via the (CAU)<sub>4</sub> and (CUA)<sub>4</sub>  
10 sequences at the ends of the primers, and transformed into *E. coli* DH5α.

The Fad2-D and Fad2-F inserts were sequenced on both strands with an ABI PRISM 310 automated sequencer (Perkin-Elmer) following the manufacturer's directions, using synthetic primers, AmpliTaq® DNA polymerase and dye terminator.

15 The Fad2-D gene was found to have an intron upstream of the ATG start codon. As expected, the coding sequence of the gene contained a G to A mutation at nucleotide 316, derived from IMC 129 (Fig. 2).

A single base transversion from G to A at nucleotide 908 was detected in the F gene sequence of the Q4275 amplified products, compared to the wild type  
20 F gene sequence (Fig. 2). This mutation changes the codon at amino acid 303 from GGA to GAA, resulting in the non-conservative substitution of a glutamic acid residue for a glycine residue (Table 3 and Fig. 3). Expression of the mutant Q4275 Fad2-F delta-12 desaturase gene in plants alters the fatty acid composition, as described hereinabove.

25

## EXAMPLE 12

### High Erucic, High Oleic Acid Rapeseed

The breeding procedure designed to produce novel fatty acid compositions in rapeseed is outlined in Figure 4. In general, crosses were made between a high erucic acid line and a high oleic acid line. The high erucic acid



- 51 -

line, designated HECO1 (sold under the trade name Hero), contains about 45.5% erucic acid (Table 23). The high oleic acid lines were designated 93GS66A-130 and 93GS34A-179 and were derived from 93GS. See, for example, Example 5 and Table 17. These lines contain about 84% oleic acid in their seed oil (Table 24).

5

TABLE 23

**Fatty Acid Composition of HECO1**

10

15

20

Fatty Acid	Weight (%)
C <sub>14:0</sub>	0.05
C <sub>16:0</sub>	3.60
C <sub>16:1</sub>	0.36
C <sub>16:0</sub>	1.66
C <sub>18:1</sub>	14.72
C <sub>18:2</sub>	10.67
C <sub>14:0</sub>	9.71
C <sub>20:0</sub>	1.36
C <sub>20:1</sub>	9.04
C <sub>20:2</sub>	0.48
C <sub>22:0</sub>	1.74
C <sub>22:1</sub>	45.45
C <sub>24:0</sub>	0.49
C <sub>24:1</sub>	0.81

TABLE 24  
Fatty Acid Composition of 93GS66A-130 and 93GS34A-179

Fatty Acid	Weight(%) of 93GS66A-130	Weight(%) of 93GS34A-179
C <sub>14:0</sub>	0.04	0.05
C <sub>16:0</sub>	3.25	3.23
C <sub>16:1</sub>	0.25	0.25
C <sub>18:0</sub>	1.60	1.94
C <sub>18:1</sub>	84.38	83.71
C <sub>18:2</sub>	2.58	3.14
C <sub>18:3</sub>	4.86	4.76
C <sub>20:0</sub>	0.56	0.65
C <sub>20:1</sub>	1.57	1.41
C <sub>20:2</sub>	0.05	0.04
C <sub>22:0</sub>	0.37	0.39
C <sub>22:1</sub>	0.06	0.03
C <sub>24:0</sub>	0.20	0.18
C <sub>24:1</sub>	0.21	0.18

The F<sub>1</sub> generations of crosses between HEC01 x 93GS66A-130, and HEC01 x 93GS34A-179, were designated 96.801 and 96.804, respectively. F<sub>1</sub> 96.801 and 96.804 plants were self-pollinated to produce F<sub>2</sub> seed. Overall, 622 random single F<sub>2</sub> seeds were analyzed for their fatty acid composition. Table 25 summarizes the average percent and standard deviation for total monounsaturated content, oleic acid, eicosenoic acid, erucic acid, total polyunsaturated and total saturated fatty acid content of these 622 seeds.

- 53 -

TABLE 25

Fatty acid	%
Total long chain monounsaturated	78.90 $\pm$ 4.07
Palmitoleate	0.28 $\pm$ 0.06
Oleic Acid	45.33 $\pm$ 9.91
Eicosenoic Acid	14.84 $\pm$ 2.84
Erucic Acid	17.97 $\pm$ 8.9
Nervonic Acid	0.48 $\pm$ 0.21
Total polyunsaturated	7.10 $\pm$ 1.05
Total saturated	13.99 $\pm$ 3.83

Analysis of this data indicate that the frequency distributions deviate from a normal distribution. The total long chain monounsaturated content frequency distribution is slightly skewed to the right (-0.0513), and the eicosenoic acid content distribution is strongly skewed to the right (-1.715). Frequency distributions for oleic acid and erucic acid content are strongly skewed to the left (0.397 and 0.177, respectively). Skewness was calculated using Lotus 1-2-3 for Windows (release 5.0).

Table 26 describes characteristics of selected populations within the total population of seeds. For example, 151 seeds had a long chain monounsaturated fatty acid content greater than 82% (Table 26, column B). Within this population, the average oleic, eicosenoic and erucic acid content was about 48%, 16%, and 19%, respectively. Total polyunsaturated fatty acid content (C18:2, C18:3, and C20:2) was about 9% and total saturated fatty acid content was less than 7%.

Forty-seven of the 622 seeds had a long chain monounsaturated content greater than 85% (Table 26, column C). The average oleic, eicosenoic and erucic acid content within these seeds was 51%, 17%, and 17%, respectively. Total saturated and total polyunsaturated fatty acids were each less than 7%.

- 54 -

Twenty-three of the seeds had an eicosenoic acid content greater than 19% (Table 26, column F). Within these seeds, the average oleic acid erucic acid content was about 44% and 19%, respectively. Total polyunsaturated fatty acids were less than 10% and total saturated fatty acids were less than 7%.

Table 26

	A	B	C	D	E	F
Total Saturated	6.76 ± 0.72	6.65 ± 0.07	6.68 ± 0.61	6.85 ± 0.98	6.85 ± 0.93	6.66 ± 0.78
C14:0	0.04 ± 0.04	0.07 ± 0.05	0.06 ± 0.03	0.07 ± 0.04	0.06 ± 0.64	0.07 ± 0.05
C16:0	3.42 ± 0.37	3.35 ± 0.34	3.28 ± 0.32	3.45 ± 0.40	3.51 ± 0.42	3.31 ± 0.33
C18:0	1.92 ± 0.33	1.93 ± 0.32	2.06 ± 0.32	1.83 ± 0.30	1.90 ± 0.30	1.93 ± 0.23
C20:0	0.77 ± 0.14	0.76 ± 0.14	0.76 ± 0.13	0.80 ± 0.13	0.76 ± 0.12	0.79 ± 0.16
C22:0	0.38 ± 0.14	0.36 ± 0.12	0.37 ± 0.11	0.42 ± 0.15	0.35 ± 0.12	0.35 ± 0.29
C24:0	0.21 ± 0.19	0.19 ± 0.14	0.20 ± 0.17	0.28 ± 0.76	0.26 ± 0.71	0.19 ± 0.24
Total Monounsaturated	82.91 ± 2.11	84.21 ± 1.64	86.21 ± 1.00	79.49 ± 4.00	80.36 ± 3.75	83.92 ± 2.43
C16:1	0.28 ± 0.05	0.27 ± 0.05	0.27 ± 0.04	0.27 ± 0.05	0.27 ± 0.05	0.26 ± 0.05
C18:1	46.45 ± 9.47	47.66 ± 9.22	51.33 ± 8.96	39.29 ± 6.21	43.55 ± 6.91	44.08 ± 2.89
C20:1	16.91 ± 2.57	16.41 ± 2.47	16.72 ± 2.40	15.56 ± 2.22	16.78 ± 1.30	19.97 ± 0.63
C22:1	19.69 ± 8.45	19.38 ± 8.25	17.39 ± 12.26	23.81 ± 5.89	19.27 ± 6.39	19.09 ± 2.32
C24:1	0.49 ± 0.19	0.48 ± 0.17	0.50 ± 0.18	0.56 ± 0.19	0.49 ± 0.21	0.52 ± 0.29
Total Polyunsaturated	10.33 ± 2.10	9.14 ± 1.71	7.11 ± 0.98	13.66 ± 3.87	12.80 ± 3.60	9.43 ± 2.43
C18:2	5.16 ± 1.5	4.36 ± 1.24	3.17 ± 0.82	7.25 ± 2.62	6.58 ± 2.40	3.96 ± 1.39
C18:3	5.02 ± 1.18	4.65 ± 1.07	3.84 ± 0.69	6.19 ± 1.61	6.02 ± 1.52	5.27 ± 1.25
C20:2	0.15 ± 0.10	0.13 ± 0.04	0.10 ± 0.04	0.22 ± 0.11	0.19 ± 0.10	0.20 ± 0.28

A=>80% total long chain monounsaturated content, n=247; B=>82% total long chain monounsaturated content, n=151; C=>85% total long chain monounsaturated content, n=47; D=>15% erucic acid, n=318; E=>15% eicosenoic acid, n=323; F=>19% eicosenoic acid, n=23

- 56 -

Fatty acid composition of selected single seeds is presented in Table 27.

V800655.334 was a single seed that had a long chain monounsaturated fatty acid content of approximately 84%. The oleic acid, eicosenoic acid and erucic acid content was 33.48%, 17.14%, and 32.23%, respectively. The total polyunsaturated  
5 fatty acid content was approximately 10%. The linoleic,  $\alpha$ -linolenic and erucic acid content was 3.54%, 6.01%, and 0.15%, respectively.

V800655.126 was a single seed that had a long chain monounsaturated fatty acid content of approximately 85% (42.67% oleic acid, 16.21% eicosenoic acid, and 25.37% erucic acid). The total polyunsaturated fatty acid content was  
10 approximately 8% (4.87% linoleic acid, 3.05%  $\alpha$ -linolenic acid, and 0.13% eicosadienoic acid).

V800654.9 was a single seed that had a long chain monounsaturated fatty acid content of 89% (51.53% oleic acid, 16.94% eicosenoic acid, and 19.24% erucic acid). The total polyunsaturated fatty acid content was approximately 8%  
15 (4.87% linoleic acid, 3.05%  $\alpha$ -linolenic acid, and 0.13% eicosadienoic acid).

Single seeds having a long chain monounsaturated fatty acid content of at least about 82% and an erucic acid content of at least about 15% were planted in a greenhouse, grown to maturity and self-pollinated. Seed ( $F_3$  generation) from each plant were harvested. A bulk seed sample from each  $F_2$  plant is analyzed for  
20 fatty acid composition.

TABLE 27

Fatty Acid Composition of Selected Single Seeds

Fatty Acid	V800655.334 Weight (%)	V800655.126 Weight (%)	V800654.9 Weight (%)
C <sub>14:0</sub>	0.07	0.05	0.03
C <sub>16:0</sub>	3.49	3.52	2.98
C <sub>16:1</sub>	0.34	0.28	0.28
C <sub>18:0</sub>	1.64	1.89	1.65
C <sub>18:1</sub>	33.48	42.67	51.53
C <sub>18:2</sub>	3.54	4.87	2.09
C <sub>18:3</sub>	6.01	3.05	3.53
C <sub>20:0</sub>	0.86	0.87	0.68
C <sub>20:1</sub>	17.14	16.21	16.94
C <sub>20:2</sub>	0.15	0.13	0.10
C <sub>22:0</sub>	0.41	0.35	0.24
C <sub>22:1</sub>	32.23	25.37	19.24
C <sub>24:0</sub>	0.12	0.13	0.14
C <sub>24:1</sub>	0.52	0.61	0.59

EXAMPLE 13

Additional crosses were made between Hero and several high oleic lines (Table 28) to increase the seed erucic acid content through a reduction in polyunsaturates content and increase in total monounsaturates content. The high oleic acid lines included 048X058 and Q4275X663-40. The 048X058 line resulted from a cross of two separate transformed lines. The 048X058 line contains a co-suppression event resulting from introduction of the 663-40 transgene described above, and a second co-suppression event resulting from a transgene that includes an oleosin promoter linked to an oleic desaturase gene. The Q4275X663-40 line was derived from a cross of Q4275 (Example 5 and Table 17) by 663-40. The

- 58 -

- 663-40 line contains a co-suppression event resulting from a transgene that includes a napin promotor linked to a linoleic desaturase gene. Plants of each line were grown in growth chambers under 16 hrs of light at 23/17°C day/night temperature. Flowers were emasculated prior to opening and covered to prevent cross
- 5 pollination. On the following day, stigmas of emasculated flowers were pollinated with the desired pollen donor. At pod maturity the F1 seed was harvested.



TABLE 28  
Crossing block

Cross Number	Female Parent	Female Phenotype	Male Parent	Male Phenotype	Source of Male Phenotype
HEHOA	HEC101	High 22:1	048X058	High 18:1/ Low 18:3	Transgenes
HEHOB	HEC101	High 22:1	Q4275X663-40	High 18:1/ Low 18:3	Mutant/ Transgene
HEHOC	HEC101	High 22:1	Q4275X663-40	High 18:1/ Low 18:3	Mutant/ Transgene

- 60 -

F1 seed generated from the crosses in Table 28 were advanced to F2 seed generation in the growth chamber. Ten seeds were individually planted for each cross. At flowering the plants were covered with bags to ensure self pollination. The F2 seeds were harvested at maturity.

5           The seeds were germinated on filter paper at room temperature in the dark. Eighteen to 24 hours after the start of germination, one cotyledon was removed from the seed for extraction of fatty acids. Fatty acid compositions were determined using gas chromatography. Selected F2 half seeds having a high erucic content are shown in Tables 29 and 30.

10           F<sub>2</sub> half seeds were planted in soil and grown under growth chamber conditions described above. At flowering the plants were covered with bags for self pollination. After maturity, F<sub>3</sub> seed was harvested and analyzed for fatty acid composition. Seeds were analyzed using a 10-15 seed sample size. The results of the analysis are in Tables 31 and 32.

TABLE 29  
Fatty Acid Composition of F<sub>1</sub> Half Seeds of HEHOA [HE101X(048X052)]

Sample No.	Fatty Acid Composition (%)										
	C16:0	C18:0	C18:1	C18:2	C18:3	C20:0	C20:1	C22:0	C22:1	C24:0	C24:1
VL10186-1	3.34	1.83	49.7	3.32	1.59	0.87	19.62	0.30	18.10	0.39	0.35
VL10186-5	2.61	1.07	29.14	5.81	2.42	0.71	14.99	0.31	40.90	0.93	0.60
VL10186-33	3.47	1.32	29.73	4.38	2.98	0.86	12.22	0.44	41.21	1.50	1.28
VL10186-59	4.01	1.68	41.38	2.96	1.69	1.08	19.72	0.43	26.05	0	0.55
VL10186-67	3.90	1.29	29.10	3.65	2.89	0.88	13.79	0.52	40.96	1.31	1.09
VL10186-74	2.76	1.25	34.04	2.63	1.45	0.75	16.64	0.38	38.67	0.14	0.95
VL10186-88	3.23	1.39	48.97	3.27	1.50	0.60	19.71	0.17	20.48	0	0.36

5

10

TABLE 30

Fatty Acid Compositions of Half Seeds of HEHOC [HE101X(Q4275X663-40)]

Sample No.	Fatty Acid Composition (%)										
	C16:0	C18:0	C18:1	C18:2	C18:3	C20:0	C20:1	C22:0	C22:1	C24:0	C24:1
VL10200-214	2.24	0.74	31.66	3.01	6.24	0.46	11.79	0.41	40.60	0.86	1.57
VL10200-231	3.89	1.03	31.51	12.50	2.41	0.54	14.17	0.29	32.23	0	0.75
VL10200-238	3.36	0.95	33.19	8.99	1.66	0.55	14.35	0.21	33.61	0.83	1.02
VL10200-267	3.12	1.02	30.18	7.61	1.52	0.59	14.53	0.19	39.41	0.24	1.013
VL10203-50	2.63	0.97	31.79	8.47	1.99	0.58	14.58	0.25	37.41	0.13	0.59
VL10200-293	2.71	0.78	32.83	6.85	1.88	0.46	13.11	0.32	39.18	0.82	0.67

TABLE 31  
Fatty acid composition of F3 lines of 97HEHOA HE101X(048X052)I

Sample No.	Fatty Acid Composition (%)										
	C16:0	C18:0	C18:1	C18:2	C18:3	C20:0	C20:1	C22:0	C22:1	C24:0	C24:1
HEHOA-74	2.51	1.03	27.44	3.72	3.55	0.65	13.60	0.30	45.57	0.14	1.03
HEHOA-01	3.66	1.40	47.79	6.47	2.97	0.62	19.23	0.25	16.11	0.14	0.76
HEHOA-67	2.47	0.84	20.43	7.25	3.89	0.69	10.33	0.47	52.09	0.16	0.86
HEHOA-59	2.81	1.08	27.01	7.88	2.82	0.69	16.15	0.32	39.68	0.15	0.86
HEHOA-88	3.16	1.21	44.85	9.21	2.31	0.49	16.30	0.21	21.00	0.12	0.58
HEHOA-33	2.53	0.79	21.90	9.52	3.55	0.53	11.51	0.31	47.59	0.13	1.08
HEHOA-5	2.93	1.01	23.67	10.26	2.00	0.63	14.34	0.38	42.98	0.15	1.06

5

10

TABLE 32  
Fatty acid composition of F3 lines of HEHOC [HE101X(Q4275X663-40)]

Sample No.	Fatty Acid Composition (%)										
	C16:0	C18:0	C18:1	C18:2	C18:3	C20:0	C20:1	C22:0	C22:1	C24:0	C24:1
HEHOC-214	2.47	1.12	31.15	3.77	3.84	0.77	13.78	0.43	41.15	0.17	0.97
HEHOC-267	2.62	1.42	31.64	6.44	1.30	0.84	15.64	0.39	38.15	0.16	0.95
HEHOC-293	2.73	1.13	32.08	7.23	2.18	0.72	14.88	0.41	37.17	0.17	0.81
HEHOC-238	2.90	1.05	35.20	9.37	1.76	0.66	14.88	0.38	32.05	0.1	1.01
HEHOC(2)-50	2.60	0.93	31.16	5.66	2.09	0.61	14.93	0.31	40.30	0.11	0.88
HEHOC(2)-156	3.19	1.71	46.56	3.05	1.59	0.94	16.41	0.40	24.67	0.19	0.83

- 65 -

HEHOC-214 F<sub>3</sub> seeds were planted and grown under growth chamber conditions and self-pollinated as described above. At maturity, F<sub>4</sub> seeds were harvested and analyzed for fatty acid composition, using bulk samples of 10-15 seeds. The seed fatty acid composition of three F<sub>4</sub> lines is shown in Table 33. All  
5 three samples had a long chain monounsaturated fatty acid content of greater than 82% and an erucic acid content of greater than 37% based on total fatty acid composition. Genes affecting fatty acid composition are still segregating in this F<sub>4</sub> generation material. Selection in subsequent generations will fix the genetic makeup and result in lines having a seed fatty acid composition of about 25-30%  
10 oleic acid, about 3-4% linoleic acid, about 1-2%  $\alpha$ -linolenic acid, about 45-50% erucic acid and about 10-13% eicosenoic acid.

TABLE 33  
Fatty acid composition of F4 lines of HEHOC 214

Sample No.	Fatty Acid Composition (%)											Total Sats
	C16:0	C18:0	C18:1	C18:2	C18:3	C20:0	C20:1	C22:0	C22:1	C24:0	C24:1	
1	1.91	1.33	35.9	3.17	1.21	0.97	16.57	0.52	37.1	0.23	0.76	4.9
2	1.86	0.87	24.26	8.49	1.01	0.7	12.41	0.4	48.56	0.18	0.90	4.0
3	1.94	0.83	26.67	8.31	1.02	0.64	14.17	0.34	44.71	0.14	0.82	3.9



- 67 -

EXAMPLE 14

Seeds from the lines of Table 31 were planted in two separate plots. Plants in each plot were allowed to open-pollinate. Oil was extracted from seeds produced on each plot; the fatty acid compositions of each oil are shown in Table

5 34. A sample of each oil was refined and bleached (high long chain monounsaturate plot 1) or refined, bleached, and deodorized (high long chain monounsaturate plot 2). Oil from plot 1 had a total saturated fatty acid content of 5.55%, a long chain monounsaturated fatty acid content of 85.23%, and a total polyunsaturated fatty acid content of 6.80%. Oil from plot 2 had a total saturated

10 fatty acid content of 5.22%, a long chain monounsaturated fatty acid content of 82.90%, and a total polyunsaturated fatty acid content of 9.56%. The iodine values of oil from plots 1 and 2 were 79 and 81.7, respectively. Oil from plot 1, which was not deodorized, contained 420 ppm of tocopherol. Oil from plot 2 contained 280 ppm of tocopherol. Oil from plots 1 and 2 had average oxidative stabilities of

15 70 AOM hours (n=2, 69 and 71 AOM hours) and 49.5 AOM hours (n=2, 48 and 51 AOM hours), respectively.

TABLE 34

Fatty Acid Composition of High Long Chain Monounsaturate Rapeseed Oil

20

Plot No.	Fatty Acid Composition (%)								
	C16:0	C18:0	C18:1	C18:2	C18:3	C20:0	C20:1	C22:0	C22:1
Plot 1	2.45	1.50	31.50	3.68	2.45	0.90	13.20	0.39	39.3
Plot 2	2.55	1.30	29.70	6.35	2.46	0.76	12.50	0.36	39.50

Table 35 provides characteristics of oils from plots 1 and 2, which were determined by differential scanning calorimetry using a Perkin Elmer Model 7

25 differential scanning calorimeter. Samples of 7-12 mg were placed in sample pans, sealed, and loaded into an autosampler. Samples were cooled from an initial temperature of 20°C, which was maintained for 1 minute, to -30°C at a rate of

- 68 -

40°C per minute. Samples were held at -30°C for 10 minutes, then heated to 75°C at a rate of 5°C per minute to obtain a melting curve. Samples were held at 75°C for 10 minutes, then cooled to -30°C at a rate of 5°C per minute to obtain a cooling curve. The results show that these oils, having a long chain

- 5 monounsaturate content of greater than 82% and an erucic acid content of greater than 15%, exhibit melting points of about 2-3° C. By comparison, trierucin is a solid at room temperature.

TABLE 35

Characteristics of High Long Chain Monounsaturate Rapeseed Oil

10

Plot	MP (°C)	Onset of Crystallization (°C)	ΔH (j/g)
1	3.2	-24	88.5
2	2.2	-27	92.6

EXAMPLE 15

- Seeds of *Brassica napus* variety IMC 129 were mutagenized with
- 15 MNNG as described in Example 5. Treated seeds were grown as described in Example 5 and selection for decreased seed stearic acid or decreased palmitic acid content was carried out at the M3 generation. Plants from two selected lines, designated ZW1441 (decreased palmitic acid) and Y30137 (decreased stearic acid), were crossed with HE101. ZW1441XHE101 progeny that produced seeds having
- 20 decreased palmitic acid and elevated erucic acid were selected. Y30137XHE101 progeny that produced seeds having decreased stearic acid and elevated erucic acid were selected. The fatty acid composition of seeds from representative F<sub>4</sub> generation selected lines is shown in Table 36. The results show that seeds having a long chain monounsaturate content of about 82% or greater, an erucic acid
- 25 content of 15% or greater and a total saturates content of less than 4% (e.g., about 2.0 to about 4.0%) can be achieved.

TABLE 36  
Fatty Acid Composition of F<sub>4</sub> Lines

Sample No.	Fatty Acid Composition (%)									
	C16:0	C18:0	C18:1	C18:2	C18:3	C20:0	C20:1	C22:0	C22:1	
ZW1441	2.73	1.74	73.46	11.46	7.52	0.67	1.47	0.34	0	
ZW1441 x HE101										
1	1.99	0.97	14.89	10.66	6.77	0.97	8.57	0.78	52.4	
2	2.15	0.90	14.89	12.09	7.01	0.80	8.84	0.58	50.7	
3	2.18	1.02	13.86	11.24	6.64	0.97	8.89	0.65	52.33	
Y30137	3.44	1.14	78.08	7.41	6.23	0.71	1.70	0.46	0.04	
HE101 x Y30137										
1	2.26	0.77	17.17	9.31	6.32	0.69	10.44	0.50	50.99	
2	1.99	0.75	26.42	8.48	7.47	0.58	12.97	0.39	39.5	
3	1.99	0.79	24.62	5.22	6.32	0.64	13.84	0.35	44.86	

5

10

- 70 -

To the extent not already indicated, it will be understood by those of ordinary skill in the art that any one of the various specific embodiments herein described and illustrated may be further modified to incorporate features shown in other of the specific embodiments. All patents, publications, and other references  
5 cited herein are incorporated by reference in their entirety.

The foregoing detailed description has been provided for a better understanding of the invention only and no unnecessary limitation should be understood therefrom as some modifications will be apparent to those skilled in the art without deviating from the spirit and scope of the appended claims.

- 71 -

## WHAT IS CLAIMED IS:

1. A *Brassica* plant producing seeds having a long chain monounsaturated fatty acid content of at least about 82% and an erucic acid content of at least about 15% based on total fatty acid composition.
- 5 2. The plant of claim 1, said seeds having an oleic acid content of at least about 37% based on total fatty acid composition.
3. The plant of claim 1, said seeds having an eicosenoic acid content of at least about 14% based on total fatty acid composition.
4. The plant of claim 1, wherein said monounsaturated fatty acid content is from  
10 about 85% to about 90%.
5. The plant of claim 4, said seeds having an oleic acid content of at least about 42% based on total fatty acid composition.
6. The plant of claim 5, wherein said oleic acid content is from about 47% to about 56%.
- 15 7. The plant of claim 4, said seeds having an erucic acid content of from about 17% to about 31% based on total fatty acid composition.
8. The plant of claim 4, said seeds having an eicosenoic acid content from about 15% to about 21% based on total fatty acid composition.
9. The plant of claim 1, said seeds having a saturated fatty acid content of less  
20 than about 7% based on total fatty acid composition.

- 72 -

10. The plant of claim 9, said seeds having a saturated fatty acid acid content of less than about 4% based on total fatty acid composition.
11. The plant of claim 10, said seeds having a saturated fatty acid acid content of about 2% to about 4% based on total fatty acid composition.
- 5 12. The plant of claim 1, said seeds having a polyunsaturated fatty acid content of less than about 11% based on total fatty acid composition.
13. The plant of claim 12, said seeds having a polyunsaturated fatty acid content of about 6% to about 11% based on total fatty acid composition.
14. The plant of claim 1, said seeds having an  $\alpha$ -linolenic acid content of about 1%  
10 to about 2% based on total fatty acid composition.
15. Progeny of the plant of claim 1, said progeny having said long chain monounsaturated fatty acid content and said erucic acid content.
16. A *Brassica* seed oil having a long chain monounsaturated fatty acid content of at least about 82% and an erucic acid content of at least about 15% based on total  
15 fatty acid composition.
17. The oil of claim 16, said oil having an oleic acid content of at least about 37% based on total fatty acid composition.
18. The oil of claim 16, said oil having an eicosenoic acid content of at least about 14% based on total fatty acid composition.
- 20 19. The oil of claim 16, wherein said monounsaturated fatty acid content is from about 85% to about 90%.

- 73 -

20. The oil of claim 19, said oil having an oleic acid content of at least about 42% based on total fatty acid composition.
21. The oil of claim 20, wherein said oleic acid content is from about 47% to about 56%.
- 5 22. The oil of claim 19, said oil having an erucic acid content of from about 17% to about 31% based on total fatty acid composition.
23. The oil of claim 19, said oil having an eicosenoic acid content from about 15% to about 21% based on total fatty acid composition.
24. The oil of claim 16, said oil having a saturated fatty acid content of less than  
10 about 7% based on total fatty acid composition.
25. The oil of claim 16, said oil having a polyunsaturated fatty acid content of less than about 11% based on total fatty acid composition.
26. The oil of claim 25, wherein said polyunsaturated fatty acid content is less than about 9%.
- 15 27. A *Brassica* seed oil having a long chain monounsaturated fatty acid content of at least about 82%, wherein the sum of the nervonic acid, erucic acid and eicosenoic acid content is from about 50% to about 66% based on total fatty acid composition.
28. The seed oil of claim 27, wherein the oleic acid content is from about 25% to  
20 about 30%.

- 74 -

29. A method of producing a *Brassica* plant, said method comprising the steps of crossing a first plant line with a second plant line and selecting progeny of said cross, wherein said first plant line has an erucic acid content of at least about 45% based on total fatty acid composition and said second plant line has an oleic acid content of at least about 84% based on total fatty acid composition, said progeny having a long chain monounsaturated fatty acid content of at least about 82% and an erucic acid content of at least about 15% based on total fatty acid composition.
30. A method of making a vegetable oil, said method comprising the steps of crushing *Brassica* seeds having a long chain monounsaturated fatty acid content of at least about 82% and an erucic acid content of at least about 15% based on total fatty acid composition, and extracting said vegetable oil from said crushed seeds.
31. The method of claim 30, further comprising the steps of refining and bleaching said oil.
32. The method of claim 31, further comprising the step of deodorizing said oil.
33. A lubricant, comprising a *Brassica* oil having a long chain monounsaturated fatty acid content of at least about 82% and an erucic acid content of at least about 15% based on total fatty acid composition, and an additive.
34. The lubricant of claim 33, wherein said additive is selected from the group consisting of an antioxidant, a rust inhibitor, a corrosion inhibitor, a pour point depressant, an anti-foam additive, a colorant and a detergent.
35. The lubricant of claim 33, wherein said additive is present in an amount from about 0.01% to about 20% by weight.



- 75 -

36. A hydraulic fluid comprising a *Brassica* oil having a long chain monounsaturated fatty acid content of at least about 82% and an erucic acid content of at least about 15% based on total fatty acid composition, and an additive.

37. The hydraulic fluid of claim 36, wherein said additive is selected from the  
5 group consisting of an antioxidant, a rust inhibitor, a corrosion inhibitor, a pour point depressant, an anti-foam additive, a colorant and a detergent.

38. The hydraulic fluid of claim 36, wherein said additive is present in an amount from about 0.01% to about 20% by weight.

C18:1 Frequencies  
for 92EF (WSGA 1A X Q0508)




Fig. 1

	10	20	30	40	
1	A T G G G T G C A G G T G G A A A T G C A A G T G T C T C C T C C C A	Fad2-D wt			
1	A T G G G T G C A G G T G G A A A T G C A A G T G T C T C C T C C C A	Fad2-D (GA316) IMC 129			
1	A T G G G T G C A G G T G G A A A T G C A A G T G T C T C C T C C C A	Fad2-F wt			
1	A T G G G T G C A G G T G G A A A T G C A A G T G T C T C C T C C C A	Fad2-F (TA515) Q508			
1	A T G G G T G C A G G T G G A A A T G C A A G T G T C T C C T C C C A	Fad2-F (GA908) Q4275			
	50	60	70	80	
41	A A A A G T C T G A A A A C C G A C A A C A T C A A G C G C G T A C C C T G C C G A	Fad2-D wt			
41	A A A A G T C T G A A A A C C G A C A A C A T C A A G C G C G T A C C C T G C C G A	Fad2-D (GA316) IMC 129			
41	A G A A G T C T G A A A A C C G A C A C C A T C A A G C G C G T A C C C T G C C G A	Fad2-F wt			
41	A G A A G T C T G A A A A C C G A C A C C A T C A A G C G C G T A C C C T G C C G A	Fad2-F (TA515) Q508			
41	A G A A G T C T G A A A A C C G A C A C C A T C A A G C G C G T A C C C T G C C G A	Fad2-F (GA908) Q4275			
	90	100	110	120	
81	G A C A C C G C C C T T C A C T G T C G G A G A A C T C A A G A A A G C A A T C	Fad2-D wt			
81	G A C A C C G C C C T T C A C T G T C G G A G A A C T C A A G A A A G C A A T C	Fad2-D (GA316) IMC 129			
81	G A C A C C G C C C T T C A C T G T C G G A G A A C T C A A G A A A G C A A T C	Fad2-F wt			
81	G A C A C C G C C C T T C A C T G T C G G A G A A C T C A A G A A A G C A A T C	Fad2-F (TA515) Q508			
81	G A C A C C G C C C T T C A C T G T C G G A G A A C T C A A G A A A G C A A T C	Fad2-F (GA908) Q4275			
	130	140	150	160	
121	C C A C C G C A C T G T T T C A A A C G C T C G A T C C C C T C G C T C T T T C T	Fad2-D wt			
121	C C A C C G C A C T G T T T C A A A C G C T C G A T C C C C T C G C T C T T T C T	Fad2-D (GA316) IMC 129			
121	C C A C C G C A C T G T T T C A A A C G C T C G A T C C C C T C G C T C T T T C T	Fad2-F wt			
121	C C A C C G C A C T G T T T C A A A C G C T C G A T C C C C T C G C T C T T T C T	Fad2-F (TA515) Q508			
121	C C A C C G C A C T G T T T C A A A C G C T C G A T C C C C T C G C T C T T T C T	Fad2-F (GA908) Q4275			

Fig. 2A

	170	180	190	200	
161	C C T A C C C T C A T C T G G G A C A T C A T C A T A G C C C T C C T G C C T T C C T A	Fad2-D wt			
161	C C T A C C C T C A T C T G G G A C A T C A T C A T A G C C C T C C T G C C T T C C T A	Fad2-D (GA316)			IMC 125
161	C C T A C C C T C A T C T G G G A C A T C A T C A T A G C C C T C C T G C C T T C C T A	Fad2-F wt			
161	C C T A C C C T C A T C T G G G A C A T C A T C A T A G C C C T C C T G C C T T C C T A	Fad2-F (TA515)			Q508
161	C C T A C C C T C A T C T G G G A C A T C A T C A T A G C C C T C C T G C C T T C C T A	Fad2-F (GA908)			Q4275
	210	220	230	240	
201	C T A C G T C G C C A C C A C T T A C T T C C C T C C T C C C C T C A C C C T	Fad2-D wt			
201	C T A C G T C G C C A C C A C T T A C T T C C C T C C T C C C C T C A C C C T	Fad2-D (GA316)			IMC 125
201	C T A C G T C G C C A C C A C T T A C T T C C C T C C T C C C C T C A C C C T	Fad2-F wt			
201	C T A C G T C G C C A C C A C T T A C T T C C C T C C T C C C C T C A C C C T	Fad2-F (TA515)			Q508
201	C T A C G T C G C C A C C A C T T A C T T C C C T C C T C C C C T C A C C C T	Fad2-F (GA908)			Q4275
	250	260	270	280	
241	C T C T C C C T A C T T C G C C C T G G C C T C T C T A C T G G G C C C T G C C C A G G	Fad2-D wt			
241	C T C T C C C T A C T T C G C C C T G G C C T C T C T A C T G G G C C C T G C C C A G G	Fad2-D (GA316)			IMC 125
241	C T C T C C C T A C T T C G C C C T G G C C T C T C T A C T G G G C C C T G C C C A G G	Fad2-F wt			
241	C T C T C C C T A C T T C G C C C T G G C C T C T C T A C T G G G C C C T G C C C A G G	Fad2-F (TA515)			Q508
241	C T C T C C C T A C T T C G C C C T G G C C T C T C T A C T G G G C C C T G C C C A G G	Fad2-F (GA908)			Q4275
	290	300	310	320	
281	G C T G C C G T C C C T A A C C G G C G T C T G G G T C A T A G C C C C A C G A G T G	Fad2-D wt			
281	G C T G C C G T C C C T A A C C G G C G T C T G G G T C A T A G C C C C A C A G T G	Fad2-D (GA316)			IMC 125
281	G G T G C C G T C C C T A A C C G G C G T C T G G G T C A T A G C C C C A C G A G T G	Fad2-F wt			
281	G G T G C C G T C C C T A A C C G G C G T C T G G G T C A T A G C C C C A C G A G T G	Fad2-F (TA515)			Q508
281	G G T G C C G T C C C T A A C C G G C G T C T G G G T C A T A G C C C C A C G A G T G	Fad2-F (GA908)			Q4275

Fig. 2B

	330	340	350	360																																
321	C	G	C	C	A	C	C	G	C	C	T	T	C	A	G	C	G	A	C	T	A	C	C	A	G	T	G	G	A	C	G	A	C	Fad2-D wt		
321	C	G	C	C	A	C	C	G	C	C	T	T	C	A	G	C	G	A	C	T	A	C	C	A	G	T	G	G	A	C	G	A	C	Fad2-D (GA316)		
321	C	G	C	C	A	C	C	G	C	C	T	T	C	A	G	C	G	A	C	T	A	C	C	A	G	T	G	G	A	C	G	A	C	Fad2-F wt		
321	C	G	C	C	A	C	C	G	C	C	T	T	C	A	G	C	G	A	C	T	A	C	C	A	G	T	G	G	A	C	G	A	C	Fad2-F (TA515)		
321	C	G	C	C	A	C	C	G	C	C	T	T	C	A	G	C	G	A	C	T	A	C	C	A	G	T	G	G	A	C	G	A	C	Fad2-F (GA908)		
321	C	G	C	C	A	C	C	G	C	C	T	T	C	A	G	C	G	A	C	T	A	C	C	A	G	T	G	G	A	C	G	A	C	Q4275		
	370	380	390	400																																
361	A	C	C	G	T	C	G	G	C	C	T	C	A	T	C	T	T	C	C	A	C	T	C	C	T	T	C	C	T	C	C	T	C	Fad2-D wt		
361	A	C	C	G	T	C	G	G	C	C	T	C	A	T	C	T	T	C	C	A	C	T	C	C	T	T	C	C	T	C	C	T	C	Fad2-D (GA316)		
361	A	C	C	G	T	C	G	G	T	C	T	C	A	T	C	T	T	C	C	A	C	T	C	C	T	T	C	C	T	C	C	T	C	Fad2-F wt		
361	A	C	C	G	T	C	G	G	T	C	T	C	A	T	C	T	T	C	C	A	C	T	C	C	T	T	C	C	T	C	C	T	C	Fad2-F (TA515)		
361	A	C	C	G	T	C	G	G	T	C	T	C	A	T	C	T	T	C	C	A	C	T	C	C	T	T	C	C	T	C	C	T	C	Fad2-F (GA908)		
	410	420	430	440																																
401	A	C	T	T	C	T	C	C	T	G	G	A	A	G	T	A	C	A	G	T	C	A	T	C	G	A	C	G	C	C	A	C	C	A	Fad2-D wt	
401	A	C	T	T	C	T	C	C	T	G	G	A	A	G	T	A	C	A	G	T	C	A	T	C	G	A	C	G	C	C	A	C	C	A	Fad2-D (GA316)	
401	A	C	T	T	C	T	C	C	T	G	G	A	A	G	T	A	C	A	G	T	C	A	T	C	G	A	C	G	C	C	A	C	C	A	Fad2-F wt	
401	A	C	T	T	C	T	C	C	T	G	G	A	A	G	T	A	C	A	G	T	C	A	T	C	G	A	C	G	C	C	A	C	C	A	Fad2-F (TA515)	
401	A	C	T	T	C	T	C	C	T	G	G	A	A	G	T	A	C	A	G	T	C	A	T	C	G	A	C	G	C	C	A	C	C	A	Fad2-F (GA908)	
	450	460	470	480																																
441	C	A	C	T	G	G	C	T	C	C	C	T	C	G	A	G	A	G	A	G	A	C	G	A	A	G	T	G	T	C	C	C	C	A	G	Fad2-D wt
441	C	A	C	T	G	G	C	T	C	C	C	T	C	G	A	G	A	G	A	G	A	C	G	A	A	G	T	G	T	C	C	C	C	A	G	Fad2-D (GA316)
441	C	A	C	T	G	G	C	T	C	C	C	T	C	G	A	G	A	G	A	G	A	C	G	A	A	G	T	G	T	C	C	C	C	A	G	Fad2-F wt
441	C	A	C	T	G	G	C	T	C	C	C	T	C	G	A	G	A	G	A	G	A	C	G	A	A	G	T	G	T	C	C	C	C	A	G	Fad2-F (TA515)
441	C	A	C	T	G	G	C	T	C	C	C	T	C	G	A	G	A	G	A	G	A	C	G	A	A	G	T	G	T	C	C	C	C	A	G	Fad2-F (GA908)
																																				Q4275

Fig. 2C

	490										500										510										520										
481	A	A	G	A	G	T	C	A	G	A	C	A	T	C	A	A	G	T	G	G	T	A	C	G	G	C	A	A	G	T	A	C	C	T	C	A	A	C	A	Fad2-D wt	
481	A	A	G	A	G	T	C	A	G	A	C	A	T	C	A	A	G	T	G	G	T	A	C	G	G	C	A	A	G	T	A	C	C	T	C	A	A	C	A	Fad2-D (GA316) IMC 129	
481	A	A	G	A	G	T	C	A	G	A	C	A	T	C	A	A	G	T	G	G	T	A	C	G	G	C	A	A	G	T	A	C	C	T	C	A	A	C	A	Fad2-F wt	
481	A	A	G	A	G	T	C	A	G	A	C	A	T	C	A	A	G	T	G	G	T	A	C	G	G	C	A	A	G	T	A	C	C	T	C	A	A	C	A	Fad2-F (TA515) Q508	
481	A	A	G	A	G	T	C	A	G	A	C	A	T	C	A	A	G	T	G	G	T	A	C	G	G	C	A	A	G	T	A	C	C	T	C	A	A	C	A	Fad2-F (GA908) Q4275	
	530										540										550										560										
521	A	C	C	C	T	T	T	G	G	G	A	C	G	C	A	C	C	G	T	G	A	T	G	T	T	A	A	C	G	G	T	T	C	A	G	T	T	C	A	C	Fad2-D wt
521	A	C	C	C	T	T	T	G	G	G	A	C	G	C	A	C	C	G	T	G	A	T	G	T	T	A	A	C	G	G	T	T	C	A	G	T	T	C	A	C	Fad2-D (GA316) IMC 129
521	A	C	C	C	T	T	T	G	G	G	A	C	G	C	A	C	C	G	T	G	A	T	G	T	T	A	A	C	G	G	T	T	C	A	G	T	T	C	A	C	Fad2-F wt
521	A	C	C	C	T	T	T	G	G	G	A	C	G	C	A	C	C	G	T	G	A	T	G	T	T	A	A	C	G	G	T	T	C	A	G	T	T	C	A	C	Fad2-F (TA515) Q508
521	A	C	C	C	T	T	T	G	G	G	A	C	G	C	A	C	C	G	T	G	A	T	G	T	T	A	A	C	G	G	T	T	C	A	G	T	T	C	A	C	Fad2-F (GA908) Q4275
	570										580										590										600										
561	T	C	T	C	G	G	C	T	G	G	C	C	T	T	T	G	T	A	C	T	T	A	G	C	C	T	T	C	A	A	C	G	T	C	T	C	G	G	G	G	Fad2-D wt
561	T	C	T	C	G	G	C	T	G	G	C	C	T	T	T	G	T	A	C	T	T	A	G	C	C	T	T	C	A	A	C	G	T	C	T	C	G	G	G	G	Fad2-D (GA316) IMC 129
561	T	C	T	C	G	G	C	T	G	G	C	C	G	T	T	G	T	A	C	T	T	A	G	C	C	T	T	C	A	A	C	G	T	C	T	C	G	G	G	A	Fad2-F wt
561	T	C	T	C	G	G	C	T	G	G	C	C	G	T	T	G	T	A	C	T	T	A	G	C	C	T	T	C	A	A	C	G	T	C	T	C	G	G	G	A	Fad2-F (TA515) Q508
561	T	C	T	C	G	G	C	T	G	G	C	C	G	T	T	G	T	A	C	T	T	A	G	C	C	T	T	C	A	A	C	G	T	C	T	C	G	G	G	A	Fad2-F (GA908) Q4275
	610										620										630										640										
601	A	G	A	C	C	T	T	A	C	G	A	C	G	G	C	G	G	C	T	T	C	G	C	T	T	G	C	C	A	T	T	T	C	C	A	C	C	C	C	A	Fad2-D wt
601	A	G	A	C	C	T	T	A	C	G	A	C	G	G	C	G	G	C	T	T	C	G	C	T	T	G	C	C	A	T	T	T	C	C	A	C	C	C	C	A	Fad2-D (GA316) IMC 129
601	A	G	A	C	C	T	T	A	C	G	A	C	G	G	C	G	G	C	T	T	C	G	C	T	T	G	C	C	A	T	T	T	C	C	A	C	C	C	C	A	Fad2-F wt
601	A	G	A	C	C	T	T	A	C	G	A	C	G	G	C	G	G	C	T	T	C	G	C	T	T	G	C	C	A	T	T	T	C	C	A	C	C	C	C	A	Fad2-F (TA515) Q508
601	A	G	A	C	C	T	T	A	C	G	A	C	G	G	C	G	G	C	T	T	C	G	C	T	T	G	C	C	A	T	T	T	C	C	A	C	C	C	C	A	Fad2-F (GA908) Q4275

Fig. 2D

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Fig. 2E

	810										820										830										840														
801	A	G	T	T	T	T	G	A	T	C	A	C	T	T	A	C	T	T	G	C	A	G	C	A	C	A	C	A	C	A	T	C	C	T	T	C	C	C	T	G	Fad2-D wt				
801	A	G	T	T	T	T	G	A	T	C	A	C	T	T	A	C	T	T	G	C	A	G	C	A	C	A	C	A	C	A	T	C	C	T	T	C	C	C	T	G	Fad2-D (GA316) IMC 125				
801	C	G	T	G	T	T	G	A	T	C	A	C	T	T	A	C	T	T	G	C	A	G	C	A	C	A	C	A	C	A	T	C	C	T	T	C	C	C	T	G	Fad2-F wt				
801	C	G	T	G	T	T	G	A	T	C	A	C	T	T	A	C	T	T	G	C	A	G	C	A	C	A	C	A	C	A	T	C	C	T	T	C	C	C	T	G	Fad2-F (TA515) Q508				
801	C	G	T	G	T	T	G	A	T	C	A	C	T	T	A	C	T	T	G	C	A	G	C	A	C	A	C	A	C	A	T	C	C	T	T	C	C	C	T	G	Fad2-F (GA908) Q4275				
	850										860										870										880														
841	C	C	T	C	A	C	T	A	T	G	A	C	T	C	G	T	C	T	G	A	G	T	G	G	A	T	T	G	G	A	T	T	G	G	T	T	A	G	G	G	A	G	Fad2-D wt		
841	C	C	T	C	A	C	T	A	T	G	A	C	T	C	G	T	C	T	G	A	G	T	G	G	A	T	T	G	G	A	T	T	G	G	T	T	A	G	G	G	A	G	Fad2-D (GA316) IMC 125		
841	C	C	T	C	A	C	T	A	C	G	A	T	T	C	G	T	C	C	G	A	G	T	G	G	A	T	T	G	G	A	T	T	G	G	T	T	A	G	G	G	A	G	Fad2-F wt		
841	C	C	T	C	A	C	T	A	C	G	A	T	T	C	G	T	C	C	G	A	G	T	G	G	A	T	T	G	G	A	T	T	G	G	T	T	A	G	G	G	A	G	Fad2-F (TA515) Q508		
841	C	C	T	C	A	C	T	A	C	G	A	T	T	C	G	T	C	C	G	A	G	T	G	G	A	T	T	G	G	A	T	T	G	G	T	T	A	G	G	G	A	G	Fad2-F (GA908) Q4275		
	890										900										910										920														
881	C	T	T	T	G	G	C	C	A	C	C	C	G	T	T	G	A	C	A	G	A	G	A	C	T	A	C	G	G	A	A	T	C	T	T	G	A	A	C	A	A	Fad2-D wt			
881	C	T	T	T	G	G	C	C	A	C	C	C	G	T	T	G	A	C	A	G	A	G	A	C	T	A	C	G	G	A	A	T	C	T	T	G	A	A	C	A	A	Fad2-D (GA316) IMC 125			
881	C	T	T	T	G	G	C	T	A	C	C	C	G	T	T	G	A	C	A	G	A	G	A	C	T	A	C	G	G	A	A	T	C	T	T	G	A	A	C	A	A	Fad2-F wt			
881	C	T	T	T	G	G	C	T	A	C	C	C	G	T	T	G	A	C	A	G	A	G	A	C	T	A	C	G	G	A	A	T	C	T	T	G	A	A	C	A	A	Fad2-F (TA515) Q508			
881	C	T	T	T	G	G	C	T	A	C	C	C	G	T	T	G	A	C	A	G	A	G	A	C	T	A	C	G	G	A	A	T	C	T	T	G	A	A	C	A	A	Fad2-F (GA908) Q4275			
	930										940										950										960														
921	G	G	T	C	T	T	C	C	A	C	A	A	T	A	T	C	A	C	G	G	A	C	A	C	G	C	A	C	G	C	A	C	G	T	G	G	C	G	C	A	T	C	A	C	Fad2-D wt
921	G	G	T	C	T	T	C	C	A	C	A	A	T	A	T	C	A	C	G	G	A	C	A	C	G	C	A	C	G	C	A	C	G	T	G	G	C	G	C	A	T	C	A	C	Fad2-D (GA316) IMC 125
921	G	G	T	C	T	T	C	C	A	C	A	A	T	A	T	A	C	C	G	A	C	A	C	G	C	A	C	G	C	A	C	G	T	G	G	C	G	C	A	T	C	A	T	Fad2-F wt	
921	G	G	T	C	T	T	C	C	A	C	A	A	T	A	T	A	C	C	G	A	C	A	C	G	C	A	C	G	C	A	C	G	T	G	G	C	G	C	A	T	C	A	T	Fad2-F (TA515) Q508	
921	G	G	T	C	T	T	C	C	A	C	A	A	T	A	T	A	C	C	G	A	C	A	C	G	C	A	C	G	C	A	C	G	T	G	G	C	G	C	A	T	C	A	T	Fad2-F (GA908) Q4275	

Fig. 2F



	970	980	990	1000	
961	C T G T T C T C G A C C A T G C C G C A T T A T C A T G C C G A T G G A A G C T A	Fad2-D wt			
961	C T G T T C T C G A C C A T G C C G C A T T A T C A T G C C G A T G G A A G C T A	Fad2-D (GA316) IMC 129			
961	C T G T T C T C G A C C A T G C C G C A T T A T C A T G C C G A T G G A A G C T A	Fad2-F wt			
961	C T G T T C T C G A C C A T G C C G C A T T A T C A T G C C G A T G G A A G C T A	Fad2-F (TA515) Q508			
961	C T G T T C T C G A C C A T G C C G C A T T A T C A T G C C G A T G G A A G C T A	Fad2-F (GA908) Q4275			

	1010	1020	1030	1040	
1001	C G A A G G C G A T A A A G C C C G A T A C T G G G A G A G T A T T A T C A G T T	Fad2-D wt			
1001	C G A A G G C G A T A A A G C C C G A T A C T G G G A G A G T A T T A T C A G T T	Fad2-D (GA316) IMC 129			
1001	C C A A G G C G A T A A A G C C C G A T A C T G G G A G A G T A T T A T C A G T T	Fad2-F wt			
1001	C C A A G G C G A T A A A G C C C G A T A C T G G G A G A G T A T T A T C A G T T	Fad2-F (TA515) Q508			
1001	C C A A G G C G A T A A A G C C C G A T A C T G G G A G A G T A T T A T C A G T T	Fad2-F (GA908) Q4275			

	1050	1060	1070	1080	
1041	C G A T G G G A C G C C G G T G G T T A A G G C C G A T G T G G A G G G A G G C G	Fad2-D wt			
1041	C G A T G G G A C G C C G G T G G T T A A G G C C G A T G T G G A G G G A G G C G	Fad2-D (GA316) IMC 129			
1041	C G A T G G G A C G C C G G T G G T T A A G G C C G A T G T G G A G G G A G G C G	Fad2-F wt			
1041	C G A T G G G A C G C C G G T G G T T A A G G C C G A T G T G G A G G G A G G C G	Fad2-F (TA515) Q508			
1041	C G A T G G G A C G C C G G T G G T T A A G G C C G A T G T G G A G G G A G G C G	Fad2-F (GA908) Q4275			

*Fig. 29*

	1090	1100	1110	1120	
1081	AAGGAGTGTA	TCATGTGGAA	ACCCGGAC	ACAGGTCAGGA	Fad2-D wt
1081	AAGGAGTGTA	TCATGTGGAA	ACCCGGAC	ACAGGTCAGGA	Fad2-D (GA316) IMC 129
1081	AAGGAGTGTA	TCATGTGGAA	ACCCGGAC	ACAGGTCAGGA	Fad2-F wt
1081	AAGGAGTGTA	TCATGTGGAA	ACCCGGAC	ACAGGTCAGGA	Fad2-F (TA515) Q508
1081	AAGGAGTGTA	TCATGTGGAA	ACCCGGAC	ACAGGTCAGGA	Fad2-F (GA908) Q4275
<hr/>					
	1130	1140	1150		
1121	AGAAAGGTTGT	TCCTGGTACA	ACAAATAAGTTATGA	Fad2-D wt	
1121	AGAAAGGTTGT	TCCTGGTACA	ACAAATAAGTTATGA	Fad2-D (GA316) IMC 129	
1121	AGAAAGGTTGT	TCCTGGTACA	ACAAATAAGTTATGA	Fad2-F wt	
1121	AGAAAGGTTGT	TCCTGGTACA	ACAAATAAGTTATGA	Fad2-F (TA515) Q508	
1121	AGAAAGGTTGT	TCCTGGTACA	ACAAATAAGTTATGA	Fad2-F (GA908) Q4275	

*Fig. 2H*

Fig. 3A

Fig. 3B

170		180	
161	Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu Gly Arg Thr Val	Fad2-D wt	
161	Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu Gly Arg Thr Val	Fad2-D (GA316)	IMC129
161	Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu Gly Arg Thr Val	Fad2-F wt	
161	Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu Gly Arg Thr Val	Fad2-F (TA515)	Q508
161	Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu Gly Arg Thr Val	Fad2-F (GA908)	Q4275
190		200	
181	Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu Tyr Leu Ala Phe Asn Val Ser Gly	Fad2-D wt	
181	Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu Tyr Leu Ala Phe Asn Val Ser Gly	Fad2-D (GA316)	IMC129
181	Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu Tyr Leu Ala Phe Asn Val Ser Gly	Fad2-F wt	
181	Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu Tyr Leu Ala Phe Asn Val Ser Gly	Fad2-F (TA515)	Q508
181	Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu Tyr Leu Ala Phe Asn Val Ser Gly	Fad2-F (GA908)	Q4275
210		220	
201	Arg Pro Tyr Asp Gly Gly Phe Ala Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp	Fad2-D wt	
201	Arg Pro Tyr Asp Gly Gly Phe Ala Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp	Fad2-D (GA316)	IMC129
201	Arg Pro Tyr Asp Gly Gly Phe Ala Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp	Fad2-F wt	
201	Arg Pro Tyr Asp Gly Gly Phe Ala Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp	Fad2-F (TA515)	Q508
201	Arg Pro Tyr Asp Gly Gly Phe Ala Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp	Fad2-F (GA908)	Q4275
230		240	
221	Arg Glu Arg Leu Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu	Fad2-D wt	
221	Arg Glu Arg Leu Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu	Fad2-D (GA316)	IMC129
221	Arg Glu Arg Leu Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu	Fad2-F wt	
221	Arg Glu Arg Leu Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu	Fad2-F (TA515)	Q508
221	Arg Glu Arg Leu Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu	Fad2-F (GA908)	Q4275

Fig. 3C

												250	260
241	Tyr Arg Tyr Ala Ala Val Gln Gly Val Ala Ser Met Val Cys Phe Tyr Gly Val Pro Leu											Fad2-D wt	
241	Tyr Arg Tyr Ala Ala Val Gln Gly Val Ala Ser Met Val Cys Phe Tyr Gly Val Pro Leu											Fad2-D (GA316)	IMC129
241	Phe Arg Tyr Ala Ala Val Gln Gly Val Ala Ser Met Val Cys Phe Tyr Gly Val Pro Leu											Fad2-F wt	
241	Phe Arg Tyr Ala Ala Val Gln Gly Val Ala Ser Met Val Cys Phe Tyr Gly Val Pro Leu											Fad2-F (TA515)	Q508
241	Phe Arg Tyr Ala Ala Val Gln Gly Val Ala Ser Met Val Cys Phe Tyr Gly Val Pro Leu											Fad2-F (GA908)	Q4275
												270	280
261	Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr Leu Gln His Thr His Pro Ser Leu											Fad2-D wt	
261	Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr Leu Gln His Thr His Pro Ser Leu											Fad2-D (GA316)	IMC129
261	Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr Leu Gln His Thr His Pro Ser Leu											Fad2-F wt	
261	Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr Leu Gln His Thr His Pro Ser Leu											Fad2-F (TA515)	Q508
261	Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr Leu Gln His Thr His Pro Ser Leu											Fad2-F (GA908)	Q4275
												290	300
281	Pro His Tyr Asp Ser Ser Ser Glu Trp Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg											Fad2-D wt	
281	Pro His Tyr Asp Ser Ser Ser Glu Trp Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg											Fad2-D (GA316)	IMC129
281	Pro His Tyr Asp Ser Ser Ser Glu Trp Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg											Fad2-F wt	
281	Pro His Tyr Asp Ser Ser Ser Glu Trp Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg											Fad2-F (TA515)	Q508
281	Pro His Tyr Asp Ser Ser Ser Glu Trp Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg											Fad2-F (GA908)	Q4275
												310	320
301	Asp Tyr Gly Ile Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His											Fad2-D wt	
301	Asp Tyr Gly Ile Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His											Fad2-D (GA316)	IMC129
301	Asp Tyr Gly Ile Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His											Fad2-F wt	
301	Asp Tyr Gly Ile Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His											Fad2-F (TA515)	Q508
301	Asp Tyr Glu Ile Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His											Fad2-F (GA908)	Q4275

Fig. 3D

	Asn	Asn	Lys	Leu	ter	
381	Asn	Asn	Lys	Leu	ter	Fad2-D wt
381	Asn	Asn	Lys	Leu	ter	Fad2-D (GA316) IMC129
381	Asn	Asn	Lys	Leu	ter	Fad2-F wt
381	Asn	Asn	Lys	Leu	ter	Fad2-F (TA515) Q508
381	Asn	Asn	Lys	Leu	ter	Fad2-F (GA908) Q4275

Fig. 33




Fig. 4



## SEQUENCE LISTING

<110> Kodali, Dharma R.  
Fan, Zhegong  
DeBonte, Lorin R.

<120> PLANTS, SEEDS AND OILS HAVING AN  
ELEVATED TOTAL MONOUNSATURATED FATTY ACID CONTENT

<130> 07148/097WO1

<150> 09/128,602

<151> 1998-08-03

<160> 18

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 1155

<212> DNA

<213> Brassica napus

<220>

<221> CDS

<222> (1)...(1152)

<221> unsure

<222> (133)...(133)

<223> Xaa = Pro or Leu

<221> unsure

<222> (194)...(194)

<223> Xaa = Leu

<221> unsure

<222> (246)...(246)

<223> Xaa = Ile or Val

<221> unsure

<222> (262)...(262)

<223> Xaa = Ile or Val

<221> unsure

<222> (345)...(345)

<223> Xaa = Tyr

&lt;400&gt; 1

atg ggt gca ggt gga aga atg caa gtg tct cct ccc tcc aaa aag tct 48  
 Met Gly Ala Gly Gly Arg Met Gln Val Ser Pro Pro Ser Lys Lys Ser  
 1 5 10 15

gaa acc gac aac atc aag cgc gta ccc tgc gag aca ccg ccc ttc act 96  
 Glu Thr Asp Asn Ile Lys Arg Val Pro Cys Glu Thr Pro Pro Phe Thr  
 20 25 30

gtc gga gaa ctc aag aaa gca atc cca ccg cac tgt ttc aaa cgc tcg 144  
 Val Gly Glu Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser  
 35 40 45

atc cct cgc tct ttc tcc tac ctc atc tgg gac atc atc ata gcc tcc 192  
 Ile Pro Arg Ser Phe Ser Tyr Leu Ile Trp Asp Ile Ile Ile Ala Ser  
 50 55 60

tgc ttc tac tac gtc gcc acc act tac ttc cct ctc ctc cct cac cct 240  
 Cys Phe Tyr Tyr Val Ala Thr Thr Tyr Phe Pro Leu Leu Pro His Pro  
 65 70 75 80

ctc tcc tac ttc gcc tgg cct ctc tac tgg gcc tgc cag ggc tgc gtc 288  
 Leu Ser Tyr Phe Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val  
 85 90 95

cta acc ggc gtc tgg gtc ata gcc cac gag tgc ggc cac cac gcc ttc 336  
 Leu Thr Gly Val Trp Val Ile Ala His Glu Cys Gly His His Ala Phe  
 100 105 110

agc gac tac cag tgg ctg gac gac acc gtc ggc ctc atc ttc cac tcc 384  
 Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser  
 115 120 125

ttc ctc ctc gtc cyt tac ttc tcc tgg aag tac agt cat cga cgc cac 432  
 Phe Leu Leu Val Xaa Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His  
 130 135 140

cat tcc aac act ggc tcc ctc gag aga gac gaa gtg ttt gtc ccc aag 480  
 His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys  
 145 150 155 160

aag aag tca gac atc aag tgg tac ggc aag tac ctc aac aac cct ttg 528  
 Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu  
 165 170 175

gga cgc acc gtg atg tta acg gtt cag ttc act ctc ggc tgg cct ttg 576  
 Gly Arg Thr Val Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu  
 180 185 190

tac ttr gcc ttc aac gtc tcg ggg aga cct tac gac ggc ggc ttc gct 624  
 Tyr Xaa Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Gly Phe Ala  
 195 200 205

tgc cat ttc cac ccc aac gct ccc atc tac aac gac cgt gag cgt ctc 672  
 Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu  
 210 215 220

cag ata tac atc tcc gac gct ggc atc ctc gcc gtc tgc tac ggt ctc 720  
 Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu  
 225 230 235 240

tac cgc tac gct gct rtc caa gga gtt gcc tcg atg gtc tgc ttc tac 768  
 Tyr Arg Tyr Ala Ala Xaa Gln Gly Val Ala Ser Met Val Cys Phe Tyr  
 245 250 255

gga gtt cct ctt ctg rtt gtc aac ggg ttc tta gtt ttg atc act tac 816  
 Gly Val Pro Leu Leu Xaa Val Asn Gly Phe Leu Val Leu Ile Thr Tyr  
 260 265 270

ttg cag cac acg cat cct tcc ctg cct cac tat gac tcg tct gag tgg 864  
 Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp  
 275 280 285

gat tgg ttg agg gga gct ttg gcc acc gtt gac aga gac tac gga atc 912  
 Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile  
 290 295 300

ttg aac aag gtc ttc cac aat atc acg gac acg cac gtg gcg cat cac 960  
 Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His  
 305 310 315 320

ctg ttc tcg acc atg ccg cat tat cat gcg atg gaa gct acg aag gcg 1008  
 Leu Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala  
 325 330 335

ata aag ccg ata ctg gga gag tat tay cag ttc gat ggg acg ccg gtg 1056  
 Ile Lys Pro Ile Leu Gly Glu Tyr Xaa Gln Phe Asp Gly Thr Pro Val  
 340 345 350

gtt aag gcg atg tgg agg gag gcg aag gag tgt atc tat gtg gaa ccg 1104  
 Val Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro  
 355 360 365

gac agg caa ggt gag aag aaa ggt gtg ttc tgg tac aac aat aag tta 1152  
 Asp Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu  
 370 375 380

tga

1155

<210> 2  
 <211> 384  
 <212> PRT  
 <213> Brassica napus  
  
 <220>  
 <221> UNSURE  
 <222> (133)...(133)  
 <223> Xaa = Pro or Leu

<221> UNSURE  
 <222> (194)...(194)  
 <223> Xaa = Leu

<221> UNSURE  
 <222> (246)...(246)  
 <223> Xaa = Ile or Val

<221> UNSURE  
 <222> (262)...(262)  
 <223> Xaa = Ile or Val

<221> UNSURE  
 <222> (345)...(345)  
 <223> Xaa = Tyr

<400> 2

Met	Gly	Ala	Gly	Gly	Arg	Met	Gln	Val	Ser	Pro	Pro	Ser	Lys	Lys	Ser
1			5				10				15				
Glu	Thr	Asp	Asn	Ile	Lys	Arg	Val	Pro	Cys	Glu	Thr	Pro	Pro	Phe	Thr
		20			25				30						
Val	Gly	Glu	Leu	Lys	Lys	Ala	Ile	Pro	Pro	His	Cys	Phe	Lys	Arg	Ser
	35			40				45							
Ile	Pro	Arg	Ser	Phe	Ser	Tyr	Leu	Ile	Trp	Asp	Ile	Ile	Ile	Ala	Ser
	50			55				60							
Cys	Phe	Tyr	Tyr	Val	Ala	Thr	Thr	Tyr	Phe	Pro	Leu	Leu	Pro	His	Pro
65			70				75				80				
Leu	Ser	Tyr	Phe	Ala	Trp	Pro	Leu	Tyr	Trp	Ala	Cys	Gln	Gly	Cys	Val
		85			90				95						
Leu	Thr	Gly	Val	Trp	Val	Ile	Ala	His	Glu	Cys	Gly	His	His	Ala	Phe
	100			105				110							
Ser	Asp	Tyr	Gln	Trp	Leu	Asp	Asp	Thr	Val	Gly	Leu	Ile	Phe	His	Ser
	115			120				125							
Phe	Leu	Leu	Val	Xaa	Tyr	Phe	Ser	Trp	Lys	Tyr	Ser	His	Arg	Arg	His

130                      135                      140  
 His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys  
 145                      150                      155                      160  
 Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu  
                     165                      170                      175  
 Gly Arg Thr Val Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu  
                     180                      185                      190  
 Tyr Xaa Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Gly Phe Ala  
                     195                      200                      205  
 Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu  
                     210                      215                      220  
 Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu  
 225                      230                      235                      240  
 Tyr Arg Tyr Ala Ala Xaa Gln Gly Val Ala Ser Met Val Cys Phe Tyr  
                     245                      250                      255  
 Gly Val Pro Leu Leu Xaa Val Asn Gly Phe Leu Val Leu Ile Thr Tyr  
                     260                      265                      270  
 Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp  
                     275                      280                      285  
 Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile  
                     290                      295                      300  
 Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His  
 305                      310                      315                      320  
 Leu Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala  
                     325                      330                      335  
 Ile Lys Pro Ile Leu Gly Glu Tyr Xaa Gln Phe Asp Gly Thr Pro Val  
                     340                      345                      350  
 Val Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro  
                     355                      360                      365  
 Asp Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu  
                     370                      375                      380

&lt;210&gt; 3

&lt;211&gt; 1155

&lt;212&gt; DNA

&lt;213&gt; Brassica napus

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)...(1152)

&lt;223&gt;

&lt;221&gt; unsure

&lt;222&gt; (133)...(133)

&lt;223&gt; Xaa = Pro or Leu

&lt;221&gt; unsure

&lt;222&gt; (194)...(194)

<223> Xaa = Leu

<221> unsure

<222> (246)...(246)

<223> Xaa = Ile or Val

<221> unsure

<222> (262)...(262)

<223> Xaa = Ile or Val

<221> unsure

<222> (345)...(345)

<223> Xaa = Tyr

<400> 3

atg ggt gca ggt gga aga atg caa gtg tct cct ccc tcc aaa aag tct	48
Met Gly Ala Gly Gly Arg Met Gln Val Ser Pro Pro Ser Lys Lys Ser	
1 5 10 15	
 gaa acc gac aac atc aag cgc gta ccc tgc gag aca ccg ccc ttc act	96
Glu Thr Asp Asn Ile Lys Arg Val Pro Cys Glu Thr Pro Pro Phe Thr	
20 25 30	
 gtc gga gaa ctc aag aaa gca atc cca ccg cac tgt ttc aaa cgc tcg	144
Val Gly Glu Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser	
35 40 45	
 atc cct cgc tct ttc tcc tac ctc atc tgg gac atc atc ata gcc tcc	192
Ile Pro Arg Ser Phe Ser Tyr Leu Ile Trp Asp Ile Ile Ile Ala Ser	
50 55 60	
 tgc ttc tac tac gtc gcc acc act tac ttc cct ctc ctc cct cac cct	240
Cys Phe Tyr Tyr Val Ala Thr Thr Tyr Phe Pro Leu Leu Pro His Pro	
65 70 75 80	
 ctc tcc tac ttc gcc tgg cct ctc tac tgg gcc tgc cag ggc tgc gtc	288
Leu Ser Tyr Phe Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val	
85 90 95	
 cta acc ggc gtc tgg gtc ata gcc cac aag tgc ggc cac cac gcc ttc	336
Leu Thr Gly Val Trp Val Ile Ala His Lys Cys Gly His His Ala Phe	
100 105 110	
 agc gac tac cag tgg ctg gac gac acc gtc ggc ctc atc ttc cac tcc	384
Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser	
115 120 125	
 ttc ctc ctc gtc cyt tac ttc tcc tgg aag tac agt cat cga cgc cac	432

Phe Leu Leu Val Xaa Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His	
130 135 140	
cat tcc aac act ggc tcc ctc gag aga gac gaa gtg ttt gtc ccc aag	480
His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys	
145 150 155 160	
aag aag tca gac atc aag tgg tac ggc aag tac ctc aac aac cct ttg	528
Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu	
165 170 175	
gga cgc acc gtg atg tta acg gtt cag ttc act ctc ggc tgg cct ttg	576
Gly Arg Thr Val Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu	
180 185 190	
tac ttr gcc ttc aac gtc tcg ggg aga cct tac gac ggc ggc ttc gct	624
Tyr Xaa Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Gly Phe Ala	
195 200 205	
tgc cat ttc cac ccc aac gct ccc atc tac aac gac cgt gag cgt ctc	672
Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu	
210 215 220	
cag ata tac atc tcc gac gct ggc atc ctc gcc gtc tgc tac ggt ctc	720
Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu	
225 230 235 240	
tac cgc tac gct gct rtc caa gga gtt gcc tcg atg gtc tgc ttc tac	768
Tyr Arg Tyr Ala Ala Xaa Gln Gly Val Ala Ser Met Val Cys Phe Tyr	
245 250 255	
gga gtt cct ctt ctg rtt gtc aac ggg ttc tta gtt ttg atc act tac	816
Gly Val Pro Leu Leu Xaa Val Asn Gly Phe Leu Val Leu Ile Thr Tyr	
260 265 270	
ttg cag cac acg cat cct tcc ctg cct cac tat gac tcg tct gag tgg	864
Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp	
275 280 285	
gat tgg ttg agg gga gct ttg gcc acc gtt gac aga gac tac gga atc	912
Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile	
290 295 300	
ttg aac aag gtc ttc cac aat atc acg gac acg cac gtg gcg cat cac	960
Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His	
305 310 315 320	
ctg ttc tcg acc atg ccg cat tat cat gcg atg gaa gct acg aag gcg	1008

Leu Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala  
 325 330 335

ata aag ccg ata ctg gga gag tat tay cag ttc gat ggg acg ccg gtg 1056  
 Ile Lys Pro Ile Leu Gly Glu Tyr Xaa Gln Phe Asp Gly Thr Pro Val  
 340 345 350

gtt aag gcg atg tgg agg gag gcg aag gag tgt atc tat gtg gaa ccg 1104  
 Val Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro  
 355 360 365

gac agg caa ggt gag aag aaa ggt gtg ttc tgg tac aac aat aag tta 1152  
 Asp Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu  
 370 375 380

tga 1155

<210> 4  
 <211> 384  
 <212> PRT  
 <213> Brassica napus  
  
 <220>  
 <221> UNSURE  
 <222> (133)...(133)  
 <223> Xaa= Pro or Leu  
  
 <221> UNSURE  
 <222> (194)...(194)  
 <223> Xaa = Leu  
  
 <221> UNSURE  
 <222> (246)...(246)  
 <223> Xaa = Ile or Val  
  
 <221> UNSURE  
 <222> (262)...(262)  
 <223> Xaa = Ile or Val  
  
 <221> UNSURE  
 <222> (345)...(345)  
 <223> Xaa = Tyr

<400> 4

Met Gly Ala Gly Gly Arg Met Gln Val Ser Pro Pro Ser Lys Lys Ser  
 1 5 10 15  
 Glu Thr Asp Asn Ile Lys Arg Val Pro Cys Glu Thr Pro Pro Phe Thr



20 25 30  
 Val Gly Glu Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser  
 35 40 45  
 Ile Pro Arg Ser Phe Ser Tyr Leu Ile Trp Asp Ile Ile Ile Ala Ser  
 50 55 60  
 Cys Phe Tyr Tyr Val Ala Thr Thr Tyr Phe Pro Leu Leu Pro His Pro  
 65 70 75 80  
 Leu Ser Tyr Phe Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val  
 85 90 95  
 Leu Thr Gly Val Trp Val Ile Ala His Lys Cys Gly His His Ala Phe  
 100 105 110  
 Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser  
 115 120 125  
 Phe Leu Leu Val Xaa Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His  
 130 135 140  
 His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys  
 145 150 155 160  
 Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu  
 165 170 175  
 Gly Arg Thr Val Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu  
 180 185 190  
 Tyr Xaa Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Gly Phe Ala  
 195 200 205  
 Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu  
 210 215 220  
 Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu  
 225 230 235 240  
 Tyr Arg Tyr Ala Ala Xaa Gln Gly Val Ala Ser Met Val Cys Phe Tyr  
 245 250 255  
 Gly Val Pro Leu Leu Xaa Val Asn Gly Phe Leu Val Leu Ile Thr Tyr  
 260 265 270  
 Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp  
 275 280 285  
 Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile  
 290 295 300  
 Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His  
 305 310 315 320  
 Leu Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala  
 325 330 335  
 Ile Lys Pro Ile Leu Gly Glu Tyr Xaa Gln Phe Asp Gly Thr Pro Val  
 340 345 350  
 Val Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro  
 355 360 365  
 Asp Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu  
 370 375 380

&lt;210&gt; 5

&lt;211&gt; 1155

10

## <212> DNA

<213> Brassica napus

 $\langle 220 \rangle$ 

<221> CDS

<222> (1)...(1152)

<400> 5

atg ggt gca ggt gga aga atg caa gtg tct cct ccc tcc aag aag tct 48  
Met Gly Ala Gly Gly Arg Met Gln Val Ser Pro Pro Ser Lys Lys Ser  
1 5 10 15

gaa acc gac acc atc aag cgc gta ccc tgc gag aca ccg ccc ttc act 96  
Glu Thr Asp Thr Ile Lys Arg Val Pro Cys Glu Thr Pro Pro Phe Thr  
20 25 30

gtc gga gaa ctc aag aaa gca atc cca cgc cac tgt ttc aaa cgc tcg 144  
Val Gly Glu Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser  
35 40 45

atc cct cgc tet ttc tcc tac ctc atc tgg gac atc atc ata gcc tcc 192  
Ile Pro Arg Ser Phe Ser Tyr Leu Ile Trp Asp Ile Ile Ile Ala Ser  
50 55 60

tgc ttc tac tac gtc gcc acc act tac ttc cct ctc ctc cct cac cct 240  
Cys Phe Tyr Tyr Val Ala Thr Thr Tyr Phe Pro Leu Leu Pro His Pro  
65 70 75 80

ctc tcc tac ttc gcc tgg cct ctc tac tgg gcc tgc caa ggg tgc gtc 288  
Leu Ser Tyr Phe Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val  
85 90 95

cta acc ggc gtc tgg gtc ata gcc cac gag tgc ggc cac cac gcc ttc 336  
Leu Thr Gly Val Trp Val Ile Ala His Glu Cys Gly His His Ala Phe  
100 105 110

agc gac tac cag tgg ctt gac gac acc gtc ggt ctc atc ttc cac tcc 384  
 Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser  
 115 120 125

ttc ctc ctc gtc cct tac ttc tcc tgg aag tac agt cat cga cgc cac 432  
Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His  
130 135 140

cat tcc aac act ggc tcc ctc gag aga gac gaa gtg ttt gtc ccc aag 480  
His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys  
145 150 155 160

aag aag tca gac atc aag tgg tac ggc aag tac ctc aac aac cct ttg 528  
 Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu  
 165 170 175

gga cgc acc gtg atg tta acg gtt cag ttc act ctc ggc tgg ccg ttg 576  
 Gly Arg Thr Val Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu  
 180 185 190

tac tta gcc ttc aac gtc tcg gga aga cct tac gac ggc ggc ttc gct 624  
 Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Gly Phe Ala  
 195 200 205

tgc cat ttc cac ccc aac gct ccc atc tac aac gac cgc gag cgt ctc 672  
 Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu  
 210 215 220

cag ata tac atc tcc gac gct ggc atc ctc gcc gtc tgc tac ggt ctc 720  
 Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu  
 225 230 235 240

ttc cgt tac gcc gcc gcg cag gga gtg gcc tcg atg gtc tgc ttc tac 768  
 Phe Arg Tyr Ala Ala Ala Gln Gly Val Ala Ser Met Val Cys Phe Tyr  
 245 250 255

gga gtc ccg ctt ctg att gtc aat ggt ttc ctc gtg ttg atc act tac 816  
 Gly Val Pro Leu Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr  
 260 265 270

ttg cag cac acg cat cct tcc ctg cct cac tac gat tcg tcc gag tgg 864  
 Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp  
 275 280 285

gat tgg ttg agg gga gct ttg gct acc gtt gac aga gac tac gga atc 912  
 Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile  
 290 295 300

ttg aac aag gtc ttc cac aat att acc gac acg cac gtg gcg cat cat 960  
 Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His  
 305 310 315 320

ctg ttc tcc acg atg ccg cat tat cac gcg atg gaa gct acc aag gcg 1008  
 Leu Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala  
 325 330 335

ata aag ccg ata ctg gga gag tat tat cag ttc gat ggg acg ccg gtg 1056  
 Ile Lys Pro Ile Leu Gly Glu Tyr Tyr Gln Phe Asp Gly Thr Pro Val  
 340 345 350

gtt aag gcg atg tgg agg gag gcg aag gag tgt atc tat gtg gaa ccg 1104  
 Val Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro  
           355                  360                  365

gac agg caa ggt gag aag aaa ggt gtg ttc tgg tac aac aat aag tta 1152  
 Asp Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu  
           370                  375                  380

tga 1155

<210> 6  
 <211> 384  
 <212> PRT  
 <213> Brassica napus

<400> 6  
 Met Gly Ala Gly Gly Arg Met Gln Val Ser Pro Pro Ser Lys Lys Ser  
   1                  5                  10                  15  
 Glu Thr Asp Thr Ile Lys Arg Val Pro Cys Glu Thr Pro Pro Phe Thr  
           20                  25                  30  
 Val Gly Glu Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser  
           35                  40                  45  
 Ile Pro Arg Ser Phe Ser Tyr Leu Ile Trp Asp Ile Ile Ile Ala Ser  
           50                  55                  60  
 Cys Phe Tyr Tyr Val Ala Thr Thr Tyr Phe Pro Leu Leu Pro His Pro  
   65                  70                  75                  80  
 Leu Ser Tyr Phe Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val  
           85                  90                  95  
 Leu Thr Gly Val Trp Val Ile Ala His Glu Cys Gly His His Ala Phe  
           100                  105                  110  
 Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser  
           115                  120                  125  
 Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His  
           130                  135                  140  
 His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys  
   145                  150                  155                  160  
 Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu  
           165                  170                  175  
 Gly Arg Thr Val Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu  
           180                  185                  190  
 Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Gly Phe Ala  
           195                  200                  205  
 Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu  
           210                  215                  220  
 Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu  
   225                  230                  235                  240  
 Phe Arg Tyr Ala Ala Ala Gln Gly Val Ala Ser Met Val Cys Phe Tyr  
           245                  250                  255

13

Gly Val Pro Leu Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr  
           260                  265                  270  
 Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp  
           275                  280                  285  
 Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile  
           290                  295                  300  
 Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His  
           305                  310                  315                  320  
 Leu Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala  
           325                  330                  335  
 Ile Lys Pro Ile Leu Gly Glu Tyr Tyr Gln Phe Asp Gly Thr Pro Val  
           340                  345                  350  
 Val Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro  
           355                  360                  365  
 Asp Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu  
           370                  375                  380

<210> 7  
 <211> 1155  
 <212> DNA  
 <213> Brassica napus

<220>  
 <221> CDS  
 <222> (1)...(1152)

<400> 7

atg ggt gca ggt gga aga atg caa gtg tct cct ccc tcc aag aag tct	48
Met Gly Ala Gly Gly Arg Met Gln Val Ser Pro Pro Ser Lys Lys Ser	
1                  5                  10                  15	
gaa acc gac acc atc aag cgc gta ccc tgc gag aca ccg ccc ttc act	96
Glu Thr Asp Thr Ile Lys Arg Val Pro Cys Glu Thr Pro Pro Phe Thr	
20                  25                  30	
gtc gga gaa ctc aag aaa gca atc cca ccg cac tgt ttc aaa cgc tcg	144
Val Gly Glu Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser	
35                  40                  45	
atc cct cgc tct ttc tcc tac ctc atc tgg gac atc atc ata gcc tcc	192
Ile Pro Arg Ser Phe Ser Tyr Leu Ile Trp Asp Ile Ile Ile Ala Ser	
50                  55                  60	
tgc ttc tac tac gtc gcc acc act tac ttc cct ctc ctc cct cac cct	240
Cys Phe Tyr Tyr Val Ala Thr Thr Tyr Phe Pro Leu Leu Pro His Pro	
65                  70                  75                  80	
ctc tcc tac ttc gcc tgg cct ctc tac tgg gcc tgc caa ggg tgc gtc	288

14

Leu Ser Tyr Phe Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val	
85 90 95	
cta acc ggc gtc tgg gtc ata gcc cac gag tgc ggc cac cac gcc ttc	336
Leu Thr Gly Val Trp Val Ile Ala His Glu Cys Gly His His Ala Phe	
100 105 110	
agc gac tac cag tgg ctt gac gac acc gtc ggt ctc atc ttc cac tcc	384
Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser	
115 120 125	
ttc ctc ctc gtc cct tac ttc tcc tgg aag tac agt cat cga cgc cac	432
Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His	
130 135 140	
cat tcc aac act ggc tcc ctc gag aga gac gaa gtg ttt gtc ccc aag	480
His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys	
145 150 155 160	
aag aag tca gac atc aag tgg tac ggc aag tac cac aac aac cct ttg	528
Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr His Asn Asn Pro Leu	
165 170 175	
gga cgc acc gtg atg tta acg gtt cag ttc act ctc ggc tgg ccg ttg	576
Gly Arg Thr Val Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu	
180 185 190	
tac tta gcc ttc aac gtc tcg gga aga cct tac gac ggc ggc ttc gct	624
Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Gly Phe Ala	
195 200 205	
tgc cat ttc cac ccc aac gct ccc atc tac aac gac cgc gag cgt ctc	672
Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu	
210 215 220	
cag ata tac atc tcc gac gct ggc atc ctc gcc gtc tgc tac ggt ctc	720
Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu	
225 230 235 240	
ttc cgt tac gcc gcc gcg cag gga gtg gcc tcg atg gtc tgc ttc tac	768
Phe Arg Tyr Ala Ala Ala Gln Gly Val Ala Ser Met Val Cys Phe Tyr	
245 250 255	
gga gtc ccg ctt ctg att gtc aat ggt ttc ctc gtg ttg atc act tac	816
Gly Val Pro Leu Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr	
260 265 270	
ttg cag cac acg cat cct tcc ctg cct cac tac gat tcg tcc gag tgg	864

15

Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp  
 275 280 285

gat tgg ttg agg gga gct ttg gct acc gtt gac aga gac tac gga atc 912  
 Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile  
 290 295 300

ttg aac aag gtc ttc cac aat att acc gac acg cac gtg gcg cat cat 960  
 Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His  
 305 310 315 320

ctg ttc tcc acg atg ccg cat tat cac gcg atg gaa gct acc aag gcg 1008  
 Leu Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala  
 325 330 335

ata aag ccg ata ctg gga gag tat tat cag ttc gat ggg acg ccg gtg 1056  
 Ile Lys Pro Ile Leu Gly Glu Tyr Tyr Gln Phe Asp Gly Thr Pro Val  
 340 345 350

gtt aag gcg atg tgg agg gag gcg aag gag tgt atc tat gtg gaa ccg 1104  
 Val Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro  
 355 360 365

gac agg caa ggt gag aag aaa ggt gtg ttc tgg tac aac aat aag tta 1152  
 Asp Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu  
 370 375 380

tga 1155

<210> 8

<211> 384

<212> PRT

<213> Brassica napus

<400> 8

Met Gly Ala Gly Gly Arg Met Gln Val Ser Pro Pro Ser Lys Lys Ser  
 1 5 10 15  
 Glu Thr Asp Thr Ile Lys Arg Val Pro Cys Glu Thr Pro Pro Phe Thr  
 20 25 30  
 Val Gly Glu Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser  
 35 40 45  
 Ile Pro Arg Ser Phe Ser Tyr Leu Ile Trp Asp Ile Ile Ile Ala Ser  
 50 55 60  
 Cys Phe Tyr Tyr Val Ala Thr Thr Tyr Phe Pro Leu Leu Pro His Pro  
 65 70 75 80  
 Leu Ser Tyr Phe Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val  
 85 90 95  
 Leu Thr Gly Val Trp Val Ile Ala His Glu Cys Gly His His Ala Phe

16

100 105 110  
 Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser  
 115 120 125  
 Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His  
 130 135 140  
 His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys  
 145 150 155 160  
 Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr His Asn Asn Pro Leu  
 165 170 175  
 Gly Arg Thr Val Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu  
 180 185 190  
 Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Gly Phe Ala  
 195 200 205  
 Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu  
 210 215 220  
 Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu  
 225 230 235 240  
 Phe Arg Tyr Ala Ala Ala Gln Gly Val Ala Ser Met Val Cys Phe Tyr  
 245 250 255  
 Gly Val Pro Leu Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr  
 260 265 270  
 Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp  
 275 280 285  
 Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile  
 290 295 300  
 Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His  
 305 310 315 320  
 Leu Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala  
 325 330 335  
 Ile Lys Pro Ile Leu Gly Glu Tyr Tyr Gln Phe Asp Gly Thr Pro Val  
 340 345 350  
 Val Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro  
 355 360 365  
 Asp Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu  
 370 375 380

&lt;210&gt; 9

&lt;211&gt; 1155

&lt;212&gt; DNA

&lt;213&gt; Brassica napus

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)...(1152)

&lt;400&gt; 9

atg ggt gca ggt gga aga atg caa gtg tct cct ccc tcc aag aag tct  
 Met Gly Ala Gly Gly Arg Met Gln Val Ser Pro Pro Ser Lys Lys Ser

48



17

1	5	10	15	
gaa acc gac acc atc aag cgc gta ccc tgc gag aca ccg ccc ttc act				96
Glu Thr Asp Thr Ile Lys Arg Val Pro Cys Glu Thr Pro Pro Phe Thr				
	20	25	30	
gtc gga gaa ctc aag aaa gca atc cca ccg cac tgt ttc aaa cgc tcg				144
Val Gly Glu Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser				
	35	40	45	
atc cct cgc tct ttc tcc tac ctc atc tgg gac atc atc ata gcc tcc				192
Ile Pro Arg Ser Phe Ser Tyr Leu Ile Trp Asp Ile Ile Ala Ser				
	50	55	60	
tgc ttc tac tac gtc gcc acc act tac ttc cct ctc ctc cct cac cct				240
Cys Phe Tyr Tyr Val Ala Thr Thr Tyr Phe Pro Leu Leu Pro His Pro				
	65	70	75	80
ctc tcc tac ttc gcc tgg cct ctc tac tgg gcc tgc caa ggg tgc gtc				288
Leu Ser Tyr Phe Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val				
	85	90	95	
cta acc ggc gtc tgg gtc ata gcc cac gag tgc ggc cac cac gcc ttc				336
Leu Thr Gly Val Trp Val Ile Ala His Glu Cys Gly His His Ala Phe				
	100	105	110	
agc gac tac cag tgg ctt gac gac acc gtc ggt ctc atc ttc cac tcc				384
Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser				
	115	120	125	
ttc ctc ctc gtc cct tac ttc tcc tgg aag tac agt cat cga cgc cac				432
Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His				
	130	135	140	
cat tcc aac act ggc tcc ctc gag aga gac gaa gtg ttt gtc ccc aag				480
His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys				
	145	150	155	160
aag aag tca gac atc aag tgg tac ggc aag tac ctc aac aac cct ttg				528
Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu				
	165	170	175	
gga cgc acc gtg atg tta acg gtt cag ttc act ctc ggc tgg cgg ttg				576
Gly Arg Thr Val Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu				
	180	185	190	
tac tta gcc ttc aac gtc tcg gga aga cct tac gac ggc ggc ttc gct				624
Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Gly Phe Ala				

18

195	200	205	
tgc cat ttc cac ccc aac gct ccc atc tac aac gac cgc gag cgt ctc			672
Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu			
210	215	220	
cag ata tac atc tcc gac gct ggc atc ctc gcc gtc tgc tac ggt ctc			720
Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu			
225	230	235	240
ttc cgt tac gcc gcc gcg cag gga gtg gcc tcg atg gtc tgc ttc tac			768
Phe Arg Tyr Ala Ala Ala Gln Gly Val Ala Ser Met Val Cys Phe Tyr			
245	250	255	
gga gtc ccg ctt ctg att gtc aat ggt ttc ctc gtg ttg atc act tac			816
Gly Val Pro Leu Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr			
260	265	270	
ttg cag cac acg cat cct tcc ctg cct cac tac gat tcg tcc gag tgg			864
Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp			
275	280	285	
gat tgg ttg agg gga gct ttg gct acc gtt gac aga gac tac gaa atc			912
Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Glu Ile			
290	295	300	
ttg aac aag gtc ttc cac aat att acc gac acg cac gtg gcg cat cat			960
Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His			
305	310	315	320
ctg ttc tcc acg atg ccg cat tat cac gcg atg gaa gct acc aag gcg			1008
Leu Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala			
325	330	335	
ata aag ccg ata ctg gga gag tat tat cag ttc gat ggg acg ccg gtg			1056
Ile Lys Pro Ile Leu Gly Glu Tyr Tyr Gln Phe Asp Gly Thr Pro Val			
340	345	350	
gtt aag gcg atg tgg agg gag gcg aag gag tgt atc tat gtg gaa ccg			1104
Val Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro			
355	360	365	
gac agg caa ggt gag aag aaa ggt gtg ttc tgg tac aac aat aag tta			1152
Asp Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu			
370	375	380	
tga			1155

<210> 10  
 <211> 384  
 <212> PRT  
 <213> Brassica napus

<400> 10

```

Met Gly Ala Gly Gly Arg Met Gln Val Ser Pro Pro Ser Lys Lys Ser
1           5           10          15
Glu Thr Asp Thr Ile Lys Arg Val Pro Cys Glu Thr Pro Pro Phe Thr
20          25          30
Val Gly Glu Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser
35          40          45
Ile Pro Arg Ser Phe Ser Tyr Leu Ile Trp Asp Ile Ile Ile Ala Ser
50          55          60
Cys Phe Tyr Tyr Val Ala Thr Thr Tyr Phe Pro Leu Leu Pro His Pro
65          70          75          80
Leu Ser Tyr Phe Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val
85          90          95
Leu Thr Gly Val Trp Val Ile Ala His Glu Cys Gly His His Ala Phe
100         105         110
Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser
115         120         125
Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His
130         135         140
His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys
145         150         155         160
Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu
165         170         175
Gly Arg Thr Val Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu
180         185         190
Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Gly Phe Ala
195         200         205
Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu
210         215         220
Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu
225         230         235         240
Phe Arg Tyr Ala Ala Ala Gln Gly Val Ala Ser Met Val Cys Phe Tyr
245         250         255
Gly Val Pro Leu Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr
260         265         270
Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp
275         280         285
Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Glu Ile
290         295         300
Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His
305         310         315         320
Leu Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala
325         330         335

```

Ile Lys Pro Ile Leu Gly Glu Tyr Tyr Gln Phe Asp Gly Thr Pro Val  
                   340                  345                  350  
 Val Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro  
                   355                  360                  365  
 Asp Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu  
                   370                  375                  380

<210> 11  
 <211> 21  
 <212> DNA  
 <213> Other nucleic acid

<400> 11  
 ggatatgatg atggtgaaag a 21

<210> 12  
 <211> 21  
 <212> DNA  
 <213> Other nucleic acid

<400> 12  
 tctttcacca tcatcatatc c 21

<210> 13  
 <211> 21  
 <212> DNA  
 <213> Other nucleic acid

<400> 13  
 gttatgaagc aaagaagaaa c 21

<210> 14  
 <211> 21  
 <212> DNA  
 <213> Other nucleic acid

<400> 14  
 gttttcttct tgcttcataa c 21

<210> 15  
 <211> 32  
 <212> DNA  
 <213> Other nucleic acid

<400> 15  
 caucaucauc aucttcttcg tagggttcat cg 32

<210> 16

21

<211> 33  
<212> DNA  
<213> Other nucleic acid

<400> 16

cuacuacuac uatcatagaa gagaaagggt cag

33

<210> 17  
<211> 32  
<212> DNA  
<213> Other

<400> 17

caucaucauc aukatgggtg cacgtggaag aa

32

<210> 18  
<211> 33  
<212> DNA  
<213> Other

<400> 18

cuacuacuac uatctttcac catcatcata tcc

33

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/17645

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) : A01H 5/00, 1/02; C09K 7/02; C12P 7/40, 7/64; C12N 9/20

US CL : 800/264, 298; 426/601; 305/117

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 800/264, 298; 426/601; 305/117

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
WEST/DERWENT, DIALOG

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	TOPFER et al. Modification of Plant Lipid Synthesis. Science. 05 May 1995, Vol. 268, pages 681-686, see entire document.	1-38
Y	AXTELL, J.D. Breeding for Improved Nutritional Quality. In: Plant Breeding II (Frey, ed.) 1981, pages 365-415, see especially pages 389-395.	1-38
Y	CARR, R. Processing of Oilseed Crops. In: Oil Crops of the World (Robbelen et al, eds). 1989, pages 226-259, see especially pages 253 and 255.	16-28, 30-38
Y	US 5,703,022 A (FLOYD) 30 December 1997, col. 1-2.	16-28, 30-38

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*B* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

25 OCTOBER 1999

Date of mailing of the international search report

24 NOV 1999

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ELIZABETH F. MCELWAIN

Telephone No. 308-0196

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/17645

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,773,391 A (LAWATE et al) 30 June 1998, col. 26-35	16-28, 30-38

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/17645**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.



# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/17645

## BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-15 and 29, drawn to a Brassica plant, progeny and a method of breeding brassica plants.

Group II, claim(s) 16-28, drawn to oil.

Group III, claim(s) 30-32, drawn to a method of making vegetable oil.

Group IV, claim(s) 33-35, drawn to a lubricant.

Group V, claim(s) 36-38, drawn to hydraulic fluid.

The inventions listed as Groups I-V do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The inventions of groups I, II, IV and V are distinct products that differ chemically, structurally and functionally, and can be made separately and used independently, and lack a corresponding special technical feature, wherein the products of groups II, IV and V can be made by alternate methods than by extraction from the plants of group I, such as by chemical synthesis, and the plants may be used in a method other than for extraction of oil, such as for vegetable production. In addition, the invention of group III is drawn to a method of making oil, and the products of groups II, IV and V may be made by alternate methods, such as by chemical synthesis, and the plants of group I can be used for different methods, such as for vegetable production. Thus the claimed inventions lack a corresponding technical feature and the claimed products and methods lack unity.